



medicina *intensiva*

<http://www.medintensiva.org/en/>



ORIGINAL ARTICLE

Usefulness of vector velocity imaging in the descending thoracic aorta

Manuel Ruiz-Bailén^{a,b,*}, Javier Hidalgo-Martín^a, Julia Manetsberger^b,
Fernando Clau-Terré^c, Javier Martínez-Gámez^d, Johannes Dagomar Lohman^e,
María Leyre Lavilla-Lerma^b, Diego Fernando Matallana-Zapata^f,
Marta Ballesteros-Barroso^a, Ricardo Rivera-Fernández^a,
Pedro José Sánchez de Toro Hernández^a, José Ángel Ramos-Cuadra^g

^a Intensive Care Unit, University Jaén Hospital, Spain

^b Department of Health Sciences, University of Jaén, Spain

^c Intensive Care Medicine and Anesthesia Service, University Hospital Vall d'Hebron, Barcelona, Spain

^d Vascular Surgery Service, University of Jaén, Spain

^e Usono BV, Eindhoven, the Netherlands

^f Intensive Care Unit, Hospital Infanta Elena Huelva, Spain

^g Intensive Care Unit, Torrecárdenas Hospital, Almería, Spain

KEYWORDS

Aorta;
Speckle tracking;
Strain;
Strain rate;
Velocity;
Displacement

Abstract

Objective: To evaluate the Speckle Tracking values in the descending thoracic aorta (DTA) of a healthy population.

Design: This is an observational study.

Setting: Hospital Universitario de Jaén.

Participants: Our study included 242 healthy people (102 women) with a mean age of 42.35 ± 16.37 years.

Interventions: Evaluation of Speckle Tracking in DTA.

Main variables of interest: Strain, velocity and displacement values in the DTA were evaluated.

Results: The DAT rotational velocity was $55.91 \pm 0.22^\circ/\text{s}$, the DTA radial velocity was $0.98 \pm 0.28 \text{ cm/s}$, the circumferential strain of the DTA was $-8.98 \pm 0.87\%$, while the circumferential strain rate was $-0.96 \pm 0.15/\text{s}$. The rotational displacement of the DTA was determined at $8.87 \pm 0.36^\circ$, while radial displacement was $0.82 \pm 0.11 \text{ mm}$. Logistic regression found an association between the rotational values of the aorta and the longitudinal strain of both ventricles. In thirty people, strain, rotational velocity and displacement were also evaluated in the apical 4-chamber views and in the parasternal long-axis view. Both projections showed similar values.

DOI of original article: <https://doi.org/10.1016/j.medin.2025.502224>

* Corresponding author.

E-mail address: ruizbailen@gmail.com (M. Ruiz-Bailén).

<https://doi.org/10.1016/j.medicine.2025.502224>

2173-5727/© 2025 Elsevier España, S.L.U. and SEMICYUC. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

Please cite this article as: M. Ruiz-Bailén, J. Hidalgo-Martín, J. Manetsberger et al., Usefulness of vector velocity imaging in the descending thoracic aorta, Medicina Intensiva, <https://doi.org/10.1016/j.medicine.2025.502224>

PALABRAS CLAVE

Aorta;
Speckle tracking;
Deformidad;
Tasa de deformidad;
Velocidad;
Desplazamiento

Conclusion: Speckle Tracking could be used to study the descending thoracic aorta, and its values could suggest or rule out the presence of cardiovascular pathology.

© 2025 Elsevier España, S.L.U. and SEMICYUC. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

Utilidad del análisis de velocidad vectorial en la exploración de la aorta torácica descendente

Resumen

Objetivo: Evaluar los valores de speckle tracking en la aorta torácica descendente (ATD) en población sana.

Diseño: Estudio observacional.

Ámbito: Hospital Universitario de Jaén.

Participantes: 242 personas sanas (102 mujeres) con una edad media de $42,35 \pm 16,37$ años.

Intervenciones: Evaluación del Speckle Tracking en la ATD.

Variables de interés principales: Se ha evaluado el valor de la deformación, sus velocidades y su desplazamiento en la ATD.

Resultados: La velocidad rotacional del ATD fue de $55,91 \pm 0,22^\circ/\text{s}$, la velocidad radial del ATD fue de $0,98 \pm 0,28 \text{ cm/s}$, la deformación circunferencial de la ATD fue de $-8,98 \pm 0,87\%$, mientras que la velocidad de deformación circunferencial fue de $-0,96 \pm 0,15/\text{s}$. El desplazamiento rotacional del ATD se determinó en $8,87 \pm 0,36^\circ$, mientras que el desplazamiento radial fue de $0,82 \pm 0,11 \text{ mm}$. La regresión logística encontró asociación entre los valores rotacionales de la aorta con el strain longitudinal de ambos ventrículos. En treinta personas, también se evaluaron la tensión, la velocidad de rotación y el desplazamiento en las proyecciones apicales de 4 cámaras y en la proyección paraesternal de eje largo. Ambas proyecciones mostraron valores similares.

Conclusiones: El Speckle Tracking podría utilizarse para estudiar la aorta torácica descendente, y sus valores podrían sugerir o descartar la presencia de patología cardiovascular.

© 2025 Elsevier España, S.L.U. y SEMICYUC. Se reservan todos los derechos, incluidos los de minería de texto y datos, entrenamiento de IA y tecnologías similares.

Introduction

Classically, cardiovascular function has been based on LVEF as well as (later) diastolic dysfunction.¹ These two functions are now considered the essential basis of heart failure. However, very little is known about the atrial functions that are currently being studied extensively²⁻⁴ and even less is known about the pathophysiology of the aorta, which is still considered only a simple conducting vehicle, although recent reports may hint at further functions.³ The aorta is a fundamental organ in cardiovascular physiology, its elasticity, structure, anatomy or age change its elasticity and can modify cardiovascular physiology and may influence condition the clinical situation. Patients with atheromatous plaques, aortic artery diseases or advanced age, naturally have lower aortic distensibility and this could translate into a worsening of the symptoms.⁷⁻⁹ The aorta is distributed in three layers, i.e., the Intima, Media and Adventitia layers. The middle layer also contains muscle cells and therefore could potentially be an indicator of muscle activity.⁵ Despite the muscular tissue of the aorta being well studied, its pathophysiology is currently unknown. However, its behavior could modify the prognosis for inducing heart failure or the prog-

nosis of certain shocks.^{6,7} Our current knowledge of the aorta is based on the study of pressures, in addition to anatomical models, while the use of artificial intelligence is now also beginning to provide new novel tools and theories. The modes of evaluation of the aorta are more limited. Although progress has been made in its knowledge, further progress is needed, since it could have more functions than being a reservoir or with only modifications of the pulse wave.^{8,9} However, at present there is no method available to reliably evaluate the aorta, and to provide comparisons between healthy or diseased patients, except when evaluating its complications, dilatation, or the existence of atheromatous plaques.¹⁰

Diagnostic imaging techniques of the aortic artery such as tissue Doppler, time velocity integral, MRI or Speckle Tracking could provide physiological information that could be reflected in diagnostic and therapeutic applicability. Speckle Tracking is an avant-garde and highly efficient echocardiographic tool that has revolutionized all cardi-ological knowledge.^{3,11-14} It is much more sensitive than 2D, tissue doppler imaging and its applicability is fundamentally defined by the ability to perform strain or strain rate measurements. However, there are also offline soft-

ware tools available -e.g., vector velocity analysis¹² -which can perform measurements of longitudinal, radial or circumferential velocities and explore longitudinal, radial or rotational displacements. Despite its unparalleled potential and variability, this technology has not yet been used in the aorta, notably to evaluate the velocities, displacement, strain or strain rate of the aorta. In Intensive Care Medicine, distributive shock often occurs, affecting the entire vascular system, but we do not have tools that can assess aortic function. The possibility that Speckle Tracking in the thoracic aorta could be performed and normal patterns detected would provide another tool, in which the next step would be to correlate it with various pathologies or different hemodynamic situations and even help in patient resuscitation. Our objective was to evaluate the possibility of finding normal patterns in DTA, evaluated in a healthy population, the radial and rotational velocities, circumferential strain, strain rate, and radial or rotational displacements by vector velocity analysis echocardiography.

Methods

Study design

This is a transversal, cross sectional study aimed at creating a reference library of Speckle Tracking values and variables in the descending thoracic aorta in the healthy population. Consequently, no intervention or patient follow-up was required nor foreseen. Only anthropometric measurements were performed to assess body surface area, heart rate, weight, height, and age.

This work was approved by the local ethics committee and funded by the Department of Health of the regional government of Andalusia, Spain. The data were stored in a de-identified database to ensure the blinding of the study.

Inclusion criteria

The study was performed between January 2018 and February 2020 and from March 2022 to May 2024, with a suspension during the Covid pandemic (2020–2022). Healthy adult volunteers, who requested the echocardiography unit of the cardiological intensive care section (CCU) to perform an echocardiography, and whose echocardiography was without abnormality, were included. In addition, the study cohort also included additional volunteers who were invited to participate in the study. Participants were between 18 and 45 years old.

Exclusion criteria

People who met any of the following conditions were excluded from the study. (1) Previous cardiovascular, neurological, or renal disease. (2) Any cardiac pathology detected during echocardiography. (3) Inadequate echocardiographic study, such as poor sonographic window, or if the descending thoracic aorta was not well visible in the parasternal long-axis projection. (4) Lack of informed consent. (5) People who have presented an episode of arrhythmia or arterial

hypertension. (6) Anyone with any known disease such as endocrinopathies or diabetes.

Image acquisition and processing

Standard transthoracic echocardiography (ETT) was performed in the supine position and was recorded in digital DICOM format. For off-line analysis purposes, we used 2D and 3D, Speckle Tracking Hybrid Velocity Vector Imaging (VVI) analysis.^{11–14} Echocardiographic studies were performed with the following echocardiographic equipment available during long-term study period: Sequoia c512, Acuson, SC 2000, Siemens Medical Systems, Mountain View, CA, USA, and Epiq CX50. Transthoracic and (TTE) transesophageal echocardiography (TEE) were performed. The echocardiographic variables studied have been determined both online and offline, using echocardiographic programs Syngo software, Siemens® U.S., compatible with all DICOM formats of the different echocardiographic equipment and Tomtec Arena (<https://www.tomtec.de/products/tomtec-arena>).

The apical 4-chamber orientation was used for the acquisition of left ventricular functional data. The descending thoracic aorta was evaluated in the parasternal long-axis projection. We zoomed in on the descending thoracic aorta, and the frame rate was set as high as possible (70–120 f/s) with multiple focal points. VVI analysis was performed with a single cardiac cycle (a single heartbeat). All images were optimized with gain, compression, and dynamic range. The studies were recorded in high quality digital format. A blind, off-line analysis was conducted using the Syngo software, Siemens 2013®. U.S and Tomtec Arena® software. Separate studies were conducted to evaluate their degree of intraobserver and interobserver agreement. Figs. 1 and 2. The mean value was taken from three measurements in which the descending thoracic aorta was seen very adequately in the parasternal long-axis projection. Images with extreme values, discordant values or in which the image marked by the VVI were not adequate were rejected as outliers.

We evaluated the usual echocardiographic parameters as recommended by the American Society of Echocardiography: quantification of LVEF by 2D and 3D, Tricuspid annular plane systolic excursion (TAPSE) and the isovolumetric acceleration (IVA) calculated as the ratio of Tissue Doppler derived peak myocardial velocity during isovolumetric contraction divided by the acceleration time. The estimation of pulmonary capillary wedge pressure (PCWP) was performed using the Nagueh¹⁵ and Kuecherer equations.¹⁶ The following were also evaluated: parameters derived from speckle tracking, such as strain (S), Strain Rate (SR), longitudinal and radial displacement, and longitudinal and radial velocities in the left and the right ventricle. In the descending thoracic aorta, we calculated fractional shortening area and the following speckle-tracking VVI values as the aorta rotational and radial velocities, the aorta circumferential strain and strain rate and the thoracic aorta rotational and radial displacement. The region of interest was manually traced. Videos 1 and 2. We only assessed systolic peaks, although diastolic peaks could also have been assessed. The projection used was the parasternal long-axis, although these parameters were also used and evaluated in the apical 4-chamber projection in 30 patients. Between




Figure 1 Altman-Bland plot.

- (a) Altman-Bland plot for descending thoracic aorta Circumferential strain.
- (b) Altman-Bland plot for descending thoracic aorta Rotational velocity.
- (c) Altman-Bland plot for descending thoracic aorta Rotational displacement.
- (d) Altman-Bland plot for descending thoracic aorta circumferential strain rate.

both projections, a concordance test has been performed to evaluate the degree of agreement between these values of the VVI speckle tracking of the descending thoracic aortic artery.

Statistical analysis

A descriptive analysis of quantitative variables was performed. The R^2 coefficients of determination were evaluated by linear regression between the cardiological parameters and the values found by Speckle Tracking of the descending thoracic aorta. Degree of agreement between the results for strain, strain rate, velocities and displacement value obtained in the parasternal long axis and the apical four-chamber planes was evaluated using a Bablok regression and a Bland-Atman plot. Intraobserver and inter-operator agreement was evaluated. Results were presented using means and standard deviations. A p value <0.05 was considered statistically significant. MedCalc® Statistical Software version 22.030 (MedCalc Software Ltd, Ostend, Belgium; <https://www.medcalc.org>; 2024) and IBM SPSS for Windows, version 30.0; SPSS Inc., Chicago, Illinois, USA were used.

Results

242 analyses with good acoustic quality in healthy people were included in this study (102 women). The patients mean age was 42.35 ± 16.37 years years, their mean height

175.88 ± 16.87 cm (men) and 163.87 ± 14.71 cm (women) and the mean weight was 71.22 ± 14.45 Kg. Descending thoracic circumferential strain, circumferential strain rate, rotational velocities and displacement were evaluated in thirty people, in the apical 4-chamber and in the parasternal long-axis view. Both projections presented significant degree of agreement. Comparing results derived from the two methodologies we considered that the apical 4-chamber provides the best echocardiographic view (window), centered and zooming in the descending thoracic aorta, although it generally yields higher values than in the parasternal window (Figs. 1 and 2) (Video 1–6). The mean LVEF was 0.55 ± 0.01 using 2D echo and 0.65 ± 0.18 for 3D echo with a median heart rate of 71.44 ± 21.11 beats per minute. Global left ventricular longitudinal strain was (-21.78 ± 5.14) (Table 1). The DTA rotational velocity was $55.91 \pm 0.22^\circ/\text{s}$, the DTA radial velocity was 0.98 ± 0.28 cm/s, the circumferential strain of the DTA was $-8.98 \pm 0.87\%$, while the circumferential strain rate was $(-0.96 \pm 0.15/\text{s})$. The rotational displacement of the DTA was determined to $8.87 \pm 0.36^\circ$, and the radial displacement was 0.82 ± 0.11 mm (Figs. 3–6).

Linear regression showed and showed association between the rotational displacement of the DTA with left ventricular longitudinal strain and right ventricular longitudinal strain, (R^2 0.43, p value 0.038 and R^2 0.28, p value 0.042 respectively) and PCWP estimated by Nagueh (R^2 0.21, p value <0.001) and Kuecherer's equation (R^2 0.23, p value <0.001).




Figure 2 Regression for aorta rotational displacement in apical and parasternal projections.

Table 1 Descriptive data of the descending thoracic aorta and both ventricles.

	N	Mean	Std. deviation
Fractional shortening area	242	0.44	0.12
Aorta rotational velocity ($^{\circ}/s$)	240	55.91	0.22
Aorta radial velocity (cm/s)	212	0.98	0.28
Aorta circumferential Strain (%)	201	-8.98	0.87
Aorta Strain rate circumferential (1/s)	204	-0.96	0.15
Aorta rotational displacement ($^{\circ}$)	240	8.87	0.36
Aorta radial displacement (mm)	192	0.82	0.11
Left ventricular ejection Fraction (2D)	240	0.55	0.01
Left ventricular ejection Fraction (3D)	185	0.65	0.18
Left ventricular longitudinal global strain	235	-21.78	5.14
Right ventricular Longitudinal global strain	228	-28.13	1.32
Left ventricular Longitudinal Strain rate	218	-1.59	0.14
Lateral right ventricular strain	215	-19.02	1.32
PCWP by Nagueh equation (mmHg)	236	8.52	1.21
PCWP by Kuecherer equation (mmHg)	235	10.22	2.35
TAPSE (mm)	240	24.32	4.38
IVA (m/s^2)	237	2.92	0.27
Right ventricular Ejection Fraction 3D	146	0.55	0.18

Rotational velocities of the DTA are associated with left ventricular longitudinal strain, and right ventricular longitudinal strain, (R^2 0.44, p value 0.021 and R^2 0.36, p value 0.02 respectively). We found no relationship between these associations when adjusted for age or body mass index.

Intraoperator agreement was adequate for all parameters evaluated in the thoracic aorta, except for circumferential strain and strain rate. Similar findings were obtained for interoperator agreement (Figs. 7 and 8).

Discussion

It has been shown that aortic artery has functions that seriously impact and alter cardiovascular physiology and clinical semiology. These functions could be affected by sex, or by aldosterone inhibition, which could decrease aortic elasticity.^{17,18} Aortic artery lesions or simply atherosclerosis can also alter its functions.^{7,19}




Figure 3 Speckle Tracking VVI analysis. Exploration of the thoracic descending aorta using Speckle Tracking in the parasternal projection.

Age plays a significant role, as it causes changes in the aorta, increasing its stiffness, diameter, taper and impedance. In general, any cardiovascular pathology can modify the behavior of the aorta, which in turn can have an impact on cardiovascular behavior. These effects of changes in aortic properties and function on left ventricular contraction have been known since 2008 as ventricular-vascular interaction.^{9,19-21}

The development of heart failure after aortic repair could confirm this reciprocal interaction.²¹ Previous reports evaluating aortic diameters and aortic strain have evidenced the improvement of aortic artery elasticity with the administration of levosimendan in animal models,²² as well as after renal transplantation in patients.²³ These interactions are currently being investigated with computational models.²⁴

The most important finding of this study is that it generates the hypothesis that Speckle Tracking values, can be measured in the normal population. However, we could ask ourselves a series of questions about our findings, the first being, what is their validity when performed with software not designed for the aorta? The second question would be what would be the possible applicability of this technique? Speckle tracking is an undeniably useful tool that has advanced our knowledge of cardiac physiology.¹¹

We used Speckle Tracking designed for the cardiac cavities and traced the borders of the thoracic aorta along the

short axis of the left ventricle, assessing the systolic peak of each parameter. Obviously, it is not a specific software to assess the aorta, but we have detected some patterns and values which obviously cannot be considered as reference values due to the low population and because it has not been performed with a specific software. Working with a tool designed for another purpose makes it difficult to validate the results. We performed the study with a single cardiac cycle with a high frame rate and zooming. Sometimes the vector velocity analysis does not recognize the edges of the thoracic aorta, so the endoaortic border must be repeated. Another important limitation in its applicability is that intraoperator and interoperator agreement is not observed for strain or circumferential strain rate, although it is observed for all other parameters. Systolic peaks and diastolic peaks could be evaluated, however only systolic peaks have been evaluated. However, if this technological tool were improved, diastolic peaks could be evaluated, and more information could be obtained.

Despite all these limitations this tool together with tissue imaging or cardiac MRI could advance our understanding of aortic function could be useful and other authors have certainly used non-specific software to explore this hypothesis.²⁵⁻³⁶ The assessment of the abdominal descending aorta time velocity integral, not evaluated in this study, could also be associated with hemodynamic values



Figure 4 Radial velocity (cm/s).

and with the perfusion status and hemodynamic situation.

The applicability of aortic evaluation by speckle tracking could be very diverse. We first wanted to evaluate the values shown by this hybrid Speckle Tracking by VVI and try to associate it with the strain of the left ventricle. Its values could contribute to determining the healthy state of the aorta and even guide health professionals when assessing anatomical aortic lesions. Interestingly, we have found that the values that best correlate are not precisely the strain, nor the strain rate; but the rotational velocities and rotational displacement, which have the greatest association with cardiological values, both left and right systolic function and curiously with the estimation of the PCWP. This could help us to better treat our patients in heart failure or even septic shock or assess their physiological response to stress. One possible application of this tool is that during resuscitation of a hypovolemic septic patient with normal diastolic and systolic functions and with an elastic aorta, the patient would have at least greatly increased rotational velocities and rotational displacement. However, in the presence of myocardial dysfunction or atherosclerosis, these values or aortic deformity would be much lower. For this reason, we understand that the values obtained may have a pattern, and that in the future this technique could be applied to the aorta, to detect heart failure semiology or even diastolic dysfunction, or simply anatomical and functional alterations

of the aorta itself. The development of a specific technology to evaluate the aorta could help, for example, to differentiate true light from false light in doubtful cases of aortic dissection or facilitate its diagnosis in untrained explorers (Videos 3 and 4). In the present study we have analyzed the baseline values of the descending thoracic aorta. Obviously, these values should be confirmed in a follow up study with software designed specifically for the aorta. Our initial data however clearly shows that the analysis of the aorta using this technology can provide us with a large amount of novel information on the aorta's pathophysiology and potentially prevent the development of acute aortic syndrome in patients susceptible to them. It is hence considered a powerful new tool taking advantage of established technology. Our future aim is consequently to evaluate whether these values change in patients with different pathologies and to try to show cut-off points through which we can assess the degree of correlation of the aorta with heart failure.

Another interesting finding is the concordance detected between parasternal and apical projections, which helps their performance. This was most visible in apical 4 chamber views, where slightly higher values are obtained than in the parasternal window, a window that we have used to obtain the data Figure of the Blant-Atman plot. We have not been able to see all the parameters in all patients, so, interestingly, the ones that are best seen with this technique are rotational displacement and velocities, which are the ones




Figure 5 Rotational displacement ($^{\circ}$).

that correlate the most with cardiological parameters. Probably the parameter we found most interesting, as it is the one that correlates most with the rest of the values, is not strain but rotational displacement, which has been evaluated by other experimental techniques and has been found to be more affected in aortic annulus lesions.³⁴

We have selected those who had a very good echocardiographic window, which could generate a systematic bias that could explain this good concordance. However, we felt more comfortable with the apical 4-chamber window, a window that seemed to us to be more efficient in drawing the endoaortic border.

We used a small sample of young patients, with non-specific software, so the findings of this study should be interpreted with caution. The software used had been designed for left ventricular parameters, so we assumed that it was suitable for measuring velocities, strain and displacement. The reference values with more specific programs could probably be different, but this study clearly shows that VVI analysis can be a new and very useful tool

to study the aorta being able to evaluate its possible areas with lower velocity, deformity or motility due to fibrosis or atherosclerosis.³⁷ Study focused on the descending aorta artery we believe that it is feasible that the ascending aorta could be studied in the parasternal long axis³⁵ and even the aortic arch in the suprasternal projection.

Despite all the limitations that this manuscript presents, our results strongly suggest that Speckle Tracking can be used in the aorta, providing a new dimension of information. Using the identified reference values, aortic pathologies or cardiovascular pathological interactions could be better understood and used to explain the semiology of heart failure. Speckle Tracking applied in intensive care medicine³⁸ appears to be an effective tool. Also, in the aorta could also be used to see the impact of the aortic prosthesis repair on the patient (Videos 3 and 4).

In summary, the use of this technology in the aorta constitutes a completely new line of research, with the aim to evaluate aortic function in a reproducible manner.




Figure 6 Rotational velocity (°/s).




Figure 7 Interobserver agreement in rotational velocity.




Figure 8 Interobserver agreement in rotational displacement.

Conclusions

Speckle Tracking could be used to study the descending thoracic aorta, and its values could suggest or rule out the presence of cardiovascular pathology.

CRedit authorship contribution statement

Ruiz-Bailén, Manuel. MD, PhD 1,2. Principal investigator, author of the manuscript, database, statistical study, author of echocardiography, Martín-Hidalgo, Javier. MD 1. co-author of the manuscript, reviewer of the manuscript, collaborate in the preparation of the manuscript, translation.

Clau Terre, Fernando co-author of the manuscript, reviewer of the manuscript, collaborate in the preparation of the manuscript, translation.

Martínez-Gómez, Javier co-author of the manuscript, reviewer of the manuscript, collaborate in the preparation of the manuscript, translation.

Ramos Cuadra, José Ángel. MD. PhD4 co-author of the manuscript, reviewer of the manuscript, collaborate in the preparation of the manuscript, translation.

Lohman, Johannes Dagomar. MD5 co-author of the manuscript, reviewer of the manuscript, collaborate in the preparation of the manuscript, translation.

Manetsberger, Julia. PhD1 co-author of the manuscript, reviewer of the manuscript, collaborate in the preparation of the manuscript, translation.

Lavilla Lerma, María Leyre. PhD 1 co-author of the manuscript, reviewer of the manuscript, collaborate in the preparation of the manuscript, translation.

Funding

PAIDI CTS 606, Andalusian Health Service Project no, "PI-0585-2012 Ecocardiografía en Medicina Crítica. Detección de la disfunción Miocárdica del Paciente Crítico e interacción entre la ecocardiografía y la ventilación mecánica".

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.medin.2025.502224>.

Declaration of competing interest

The authors declare no conflict of interest related to this article.

References

1. Abdin A, Wilkinson C, Aktaa S, Böhm M, Polovina M, Rosano G, et al. European Society of Cardiology quality indicators update for the care and outcomes of adults with heart failure. The Heart Failure Association of the ESC. Eur J Heart Fail. 2024; <http://dx.doi.org/10.1002/ehf.3376>. Epub ahead of print. PMID: 38995217.
2. Tu Y, Liu X, Li X, Xue N. Left atrial stiffness index - an early marker of left ventricular diastolic dysfunction in patients with coronary heart disease. BMC Cardiovasc Disord. 2024;24(1):371, <http://dx.doi.org/10.1186/s12872-024-04047-y>. PMID: 39020270; PMCID: PMC11253388.
3. Pathan F, D'Elia N, Nolan MT, Marwick TH, Negishi K. Normal ranges of left atrial strain by speckle-tracking echocardiography.

- graphy: a systematic review and meta-analysis. *J Am Soc Echocardiogr.* 2017;30(1):59–70.e8.
4. Cesna S, Bielinis A, Zvirblis T, Miglinas M, Tarutis V. Influence of aortic arch morphology on renal perfusion in patients with coarctation of the aorta: an exploratory study. *Medicina* (Kaunas). 2024;60(6):886, <http://dx.doi.org/10.3390/medicina60060886>. PMID: 38929503; PMCID: PMC11205822.
 5. Tucker WD, Arora Y, Mahajan K. Anatomy, Blood Vessels. 2023 Aug 8. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024. PMID: 29262226.
 6. Favero AM, Rosales TO, Scheschowitsch K, Gonçalves MC, Benedet PO, Sordi R, et al. Blockade of sympathetic ganglia improves vascular dysfunction in septic shock. *Naunyn Schmiedeberg Arch Pharmacol.* 2024, <http://dx.doi.org/10.1007/s00210-024-03032-8>. Epub ahead of print. PMID: 38457039.
 7. Bell V, McCabe EL, Larson MG, Rong J, Merz AA, Osypiuk E. Relations between aortic stiffness and left ventricular mechanical function in the community. *J Am Heart Assoc.* 2017;6(1).
 8. Kanagarajan D, Heinsar S, Dau VT, Pauls JP, Tansley GD, Fraser JF. In Silico Analysis of Pulsatile Flow Veno-Arterial Extracorporeal Membrane Oxygenation on Human Aorta Model. *ASAIO J.* 2025, <http://dx.doi.org/10.1097/MAT.0000000000002418>. Epub ahead of print. PMID: 40105055.
 9. Borlaug BA, Kass DA. Ventricular-vascular interaction in heart failure. *Heart Fail Clin.* 2008;4(1):23–36, <http://dx.doi.org/10.1016/j.hfc.2007.10.001>. PMID: 18313622; PMCID: PMC2586173.4.
 10. Brie DA, Jianu AM, Popescu R, Cerbulescu T, Dema S, Brie DM, et al. Aortic aneurysms in patients with atherosclerotic coronary artery disease in the southwestern region of Romania - clinical and histopathological study. *Rom J Morphol Embryol.* 2024;65(2):251–6, <http://dx.doi.org/10.47162/RJME.65.2.11>. PMID: 39020539.
 11. Ruiz-Bailén M, Cobo Molinos J, Espada-Fuentes JC, Castillo-Rivera AM, Martínez-Amat A, Martínez Ramírez MJ. Left ventricular function improve after bench press: a speckle tracking and 3D echocardiography study. *J Sports Med Doping Stud.* 2017;7:1.
 12. Badran HM, Faheem N, Soliman M, Hamdy M, Yacoub M. Comparison of vector velocity imaging and three-dimensional speckle tracking echocardiography for assessment of left ventricular longitudinal strain in hypertrophic cardiomyopathy. *Glob Cardiol Sci Pract.* 2019;2019(1):6, <http://dx.doi.org/10.21542/gcsp.2019.6>. PMID: 31024948; PMCID: PMC6472697.
 13. Ruiz-Bailén M, Cobo-Molinos J, Espada-Fuentes JC, Castillo-Rivera AM, Martínez-Ramírez MJ, Cárdenas-Cruz A. The acute asthma: it's just a respiratory disease? A speckle tracking echocardiography study. *J Clin Respir Dis Care.* 2016;2:4, <http://dx.doi.org/10.4172/2472-1247.1000122>.
 14. Ruiz-Bailén M, Cobo-Molinos J. Myocardial dysfunction after esophageal surgery: current research. *Surgery Curr Res.* 2016;6:6, <http://dx.doi.org/10.4172/2161-1076.1000280>.
 15. Nagueh SF, Middleton KJ, Kopelen HA, Zoghbi WA, Quiñones MA. Doppler tissue imaging: a noninvasive technique for evaluation of left ventricular relaxation and estimation of filling pressures. *J Am Coll Cardiol.* 1997;30(6):1527–33, [http://dx.doi.org/10.1016/s0735-1097\(97\)00344-6](http://dx.doi.org/10.1016/s0735-1097(97)00344-6). PMID: 9362412.
 16. Kuecherer HF, Kusumoto F, Muhiudeen IA, Cahalan MK, Schiller NB. Pulmonary venous flow patterns by transesophageal pulsed Doppler echocardiography: relation to parameters of left ventricular systolic and diastolic function. *Am Heart J.* 1991;122(6):1683–93, [http://dx.doi.org/10.1016/0002-8703\(91\)90287-r](http://dx.doi.org/10.1016/0002-8703(91)90287-r). PMID: 1957763.
 17. Sakamuri A, Visniauskas B, Kilanowski-Doroh I, McNally AB, Imulinde A, Kamau A, et al. Testosterone deficiency promotes arterial stiffening independent of sex chromosome complement. *Biol Sex Differ.* 2024;15(1):46, <http://dx.doi.org/10.1186/s13293-024-00624-0>. PMID: 38845040; PMCID: PMC11155160.
 18. Vizzardi E, Pina P, Caretta G, Bonadei I, Sciatti E, Lombardi C. The effect of aldosterone-antagonist therapy on aortic elastic properties in patients with nonischemic dilated cardiomyopathy. *J Cardiovasc Med.* 2015;16:597–602.
 19. Vrublevsky AV, Boshchenko AA, Bogdanov YI, Saushkin VV, Shnaider OL, Russian, English Structural and functional disturbances of the thoracic descending aorta in atherosclerosis of various gradations. *Kardiologiya.* 2023;63(11):64–72, <http://dx.doi.org/10.18087/cardio.2023.11.n2315>. PMID: 38088114.
 20. Minderhoud SCS, van Montfoort R, Meijts TA, Korteland SA, Bruse JL, Kardys I, et al. Aortic geometry and long-term outcome in patients with a repaired coarctation. *Open Heart.* 2024;11(1):e002642, <http://dx.doi.org/10.1136/openhrt-2024-002642>. PMID: 38806222; PMCID: PMC11138275.
 21. Stonko DP, Edwards J, Abdou H, Treffalls RN, Walker P, DeMartino RR, et al. Thoracic descendent endovascular aortic repair acutely augments left ventricular biomechanics in an animal model: a mechanism for postoperative heart failure and hypertension. *Ann Vasc Surg.* 2023;97:18–26, <http://dx.doi.org/10.1016/j.avsg.2023.04.007>. Epub 2023 Apr 15. PMID: 37068623; PMCID: PMC10754260.
 22. Yang CH, Qiu HQ, Wang C, Tang YT, Zhang CR, Fan YY, et al. Levosimendan relaxes thoracic descendent aortic smooth muscle in mice by inhibiting PKC and activating inwardly rectifying potassium channels. *J Cardiovasc Pharmacol.* 2023, <http://dx.doi.org/10.1097/FJC.0000000000001524>. Epub ahead of print. PMID: 38113918.
 23. Zapolski T, Fumaga J, Jaroszynski A, Wysocka A, Rudzki S, Wysokinski A. The reverse remodeling of the aorta in patients after renal transplantation – the value of aortic stiffness index: prospective echocardiographic study. *BMC Nephrology.* 2017;18:33.
 24. Moulton MJ, Secomb TW. A fast computational model for circulatory dynamics: effects of left ventricle-aorta coupling. *Biomech Model Mechanobiol.* 2023;22(3):947–59, <http://dx.doi.org/10.1007/s10237-023-01690-w>. Epub 2023 Jan 13. PMID: 36639560; PMCID: PMC10167185.
 25. Nemes A, Domsik P, Kalapos A, Forster T. An alternative way to assess aortic elasticity: three-dimensional speckle-tracking echocardiography-derived strain assessment (From the MAGYAR-Healthy Study). *Int J Cardiol.* 2016;203:109–10, <http://dx.doi.org/10.1016/j.ijcard.2015.10.042>. Epub 2015 Oct 22. PMID: 26519670.
 26. Goyal A, Parthiban A, White DA, Forsha D, Aly D. Two-dimensional speckle tracking of the ascending aorta: a novel approach to evaluate arterial stiffness in pediatric patients with repaired conotruncal anomalies using echocardiography. *Front Cardiovasc Med.* 2025;12:1555568, <http://dx.doi.org/10.3389/fcvm.2025.1555568>. PMID: 40182427; PMCID: PMC11965654.
 27. Ruisch J, de Bakker JMK, van Helvert M, Schoonbrood MJP, Groot Jebbink E, Holeyijn S, et al. Saris AECM Ultrasound-based Velocity Vector Imaging in the Carotid Bifurcation: Repeatability and an In Vivo Comparison With 4-D Flow MRI. *Ultrasound Med Biol.* 2025;51(6):969–76.
 28. Cantinotti M, Marchese P, Scalese M, Giordano R, Franchi E, Assanta N, et al. Characterization of aortic flow patterns by high-frame-rate blood speckle tracking echocardiography in children. *J Am Heart Assoc.* 2023;12(8):e026335,

- <http://dx.doi.org/10.1161/JAHA.122.026335>. Epub 2023 Apr 17 PMID: 37066781; PMCID: PMC10227241.
29. Li Y, Deng YB, Bi XJ, Liu YN, Zhang J, Li L, et al. Evaluation of myocardial strain and aortic elasticity in patients with bicuspid aortic valve. *J Huazhong Univ Sci Technol Med Sci*. 2016;5:747–51.
30. Bieseveciene M, Vaskelyte JJ, Mizariene V, Karaliute R, Lesauskaite V, Verseckaite R. Two-dimensional speckle-tracking echocardiography for evaluation of dilative ascending aorta biomechanics. *BMC Cardiovasc Disord*. 2017;17(1):27.
31. Petrini J, Eriksson MJ, Caidahl K, Larsson M. Circumferential strain by velocity vector imaging and speckle-tracking echocardiography: validation against sonomicrometry in an aortic phantom. *Clin Physiol Funct Imaging*. 2017, <http://dx.doi.org/10.1111/cpf.12410>.
32. VIRTUE Registry Investigators. Mid-term outcomes and aortic remodelling after thoracic descendent endovascular repair for acute, subacute, and chronic aortic dissection: the VIRTUE Registry. *Eur J Vasc Endovasc Surg*. 2014;48(4):363–71.
33. Alreshidan M, Shahmansouri N, Chung J, Lash V, Emmott A, Leask R. Obtaining the biomechanical behavior of ascending aortic aneurysm via the use of novel speckle tracking echocardiography. *J Thorac Cardiovasc Surg*. 2017;153(4):781–8.
34. Kim T, Tjahjedi NS, He X, van Herwaarden JA, Patel HJ, Burris NS, et al. Three-dimensional characterization of aortic root motion by vascular deformation mapping. *J Clin Med*. 2023;12(13):4471, <http://dx.doi.org/10.3390/jcm12134471>. PMID: 37445507; PMCID.
35. Raisi-Estabragh Z, Harvey NC, Neubauer S, Petersen SE. Cardiovascular magnetic resonance imaging in the UK Biobank: a major international health research resource. *Eur Heart J Cardiovasc Imaging*. 2021;22(3):251–8, <http://dx.doi.org/10.1093/ehjci/jeaa297>. PMID: 33164079; PMCID: PMC7899275.
36. Bu Z, Ma J, Fan Y, Qiao Z, Kang Y, Zheng Y, et al. Ascending aortic strain analysis using 2-dimensional speckle tracking echocardiography improves the diagnostics for coronary artery stenosis in patients with suspected stable angina pectoris. *J Am Heart Assoc*. 2018;7(14):e008802, <http://dx.doi.org/10.1161/JAHA.118.008802>. PMID: 29982229; PMCID: PMC6064841.; PMC1034295.
37. Knopp T, Jung R, Wild J, Bochenek ML, Efentakis P, Lehmann A, et al. Myeloid cell-derived interleukin-6 induces vascular dysfunction and vascular and systemic inflammation. *Eur Heart J Open*. 2024;4(4):oeae046, <http://dx.doi.org/10.1093/ehjopen/oeae046>. PMID: 39015379; PMCID: PMC11250217.
38. Ruiz Bailén M, Hidalgo Martín J, Ramos Cuadra JA, Manetsberger J, Pola Gallego de Guzmán MD, Díaz-Castellanos MA, et al. Right ventricular strain and outflow tract velocity time integral are associated with mortality in Critically Ill patients with pulmonary embolism. *Am J Respir Crit Care Med*. 2025, <http://dx.doi.org/10.1164/rccm.202407-1433OC>.