



UPDATE IN INTENSIVE CARE MEDICINE: ULTRASOUND IN THE CRITICALLY ILL PATIENT. CLINICAL APPLICATIONS

Ultrasonography and procedures in intensive care medicine*



Pablo Blanco^{a,*}, Anselmo Abdo-Cuza^b, Elena Abril Palomares^c,
Cristina Martínez Díaz^d, Virginia Fraile Gutiérrez^e

^a High Dependency Unit, Hospital "Dr. Emilio Ferreyra", Necochea, Argentina

^b Centro de Investigaciones Médico-Quirúrgicas, La Habana, Cuba

^c Intensive Care Unit, Hospital Universitario "Torrejón de Ardoz", Madrid, Spain

^d Intensive Care Unit, Hospital Universitario "Príncipe de Asturias Alcalá de Henares", Madrid, Spain

^e Intensive Care Unit, Hospital Universitario "Rio Hortega", Valladolid, Spain

Received 2 March 2023; accepted 20 May 2023

KEYWORDS

Interventional
ultrasound;
Point-of-care;
Critical care

Abstract The use of point-of-care ultrasonography (POCUS) is not limited to the diagnosis and/or monitoring of critically ill patients. Further, ultrasound guidance is of paramount relevance to aid in successfully and safely performing several procedures in the intensive care unit (ICU). In this article, we review the role of POCUS as a procedural guidance in the ICU. Core procedures include, but are not limited to, vascular cannulation, pericardiocentesis, thoracentesis, paracentesis, aspiration of soft-tissue collections/arthrocentesis and lumbar puncture. With time, the procedures performed by intensivists may extend beyond the core competencies depicted in this review. Ultrasound guidance should be part of the intensivist's competencies, for which appropriate training should be achieved.

© 2023 Elsevier España, S.L.U. and SEMICYUC. All rights reserved.

PALABRAS CLAVE

Ecografía
intervencionista;
Sistemas de Atención
de Punto;
Cuidados críticos

Ecografía y procedimientos en Medicina Intensiva

Resumen El uso de la ecografía no se limita al diagnóstico y/o monitorización del paciente crítico. Además, la guía ecográfica es de gran importancia para la ejecución exitosa y segura de diversos procedimientos en la Unidad de Cuidados Intensivos (UCI). En este artículo, revisamos el papel de la ecografía como guía intervencionista en la UCI. Los procedimientos ecodirigidos claves son, aunque no se limitan a estos, la canulación vascular, pericardiocentesis, toracocentesis, paracentesis, aspiración de colecciones de partes blandas/artrocentesis y la punción lumbar. Con el tiempo, los procedimientos realizados por los intensivistas se extenderán más allá de los descritos en esta revisión. La guía ecográfica debe formar parte de las competencias del intensivista, para lo cual se requiere un entrenamiento adecuado.

© 2023 Elsevier España, S.L.U. y SEMICYUC. Todos los derechos reservados.

* Please cite this article as: Blanco P, Abdo-Cuza A, Palomares EA, Díaz CM, Gutiérrez VF. Ecografía y procedimientos en Medicina Intensiva. *Med Intensiva*. 2023. <https://doi.org/10.1016/j.medin.2023.05.008>

* Corresponding author.

E-mail address: ohsusabes@gmail.com (P. Blanco).

Table 1 Ultrasound-guided procedural applications in critical care patients.

Cardiovascular	Vascular cannulation* Pericardiocentesis* Temporary transvenous pacing Inferior vena cava filter placement Retrograde internal jugular vein catheterization for jugular bulb oxygen saturation monitoring
Airway, lung and pleura	Thoracentesis* Tracheal intubation Percutaneous tracheostomy Lung abscess drainage Pneumothorax drainage
Abdomen and pelvis	Paracentesis* Percutaneous cholecystostomy Organ-specific abscesses (e.g., hepatic, renal, pancreatic) and peritoneal abscess (e.g., subphrenic) Nasogastric tube placement Percutaneous gastrostomy Percutaneous nephrostomy Percutaneous cystostomy
Renal and urinary parts	Puncture of a non-deflating Foley balloon Joint aspiration* Aspiration and drainage of soft-tissue fluid collections (e.g., hematomas, abscess)*
Nervous system	Lumbar puncture* Nerve or plexus blockage

Asterisks indicates the core procedures for intensivists.

Introduction

Point-of-care ultrasonography (POCUS) has revolutionized the care of critically ill patients around the world.¹ In addition to its well-known diagnostic applications, POCUS is gaining ground as a guide for the majority of interventional procedures performed in the Intensive Care Unit (ICU).²⁻⁴

The benefits of POCUS include its fastness, absence of ionizing radiation, low cost, and bedside use without the need to move the patient to the radiology department or the operating room. The main disadvantages include operator dependency (i.e., lack of expertise) and patient dependency (i.e., inadequate ultrasound views).

In this article, we review the role of POCUS as a procedural guidance in the ICU. A list of ultrasound-guided procedural applications is provided in [Table 1](#). The core competencies of intensivists include, but are not limited to, vascular cannulation, pericardiocentesis, thoracentesis, paracentesis, soft-tissue collection aspiration/arthrocentesis, and lumbar puncture. These are described here in detail, while suggested readings are provided in [Supplementary Material, Appendix A](#), for less frequent, evolving interventions or those performed by the interventional radiologist. The general aspects of ultrasound-guided procedures are detailed in [Supplementary Material, Appendix B](#).

Vascular access

Central venous catheters (CVC) placement is the most common procedure performed under ultrasound guidance in the ICU.⁵ However, data show that the use of POCUS is suboptimal, since as many as 50% of CVC are still placed using anatomical landmarks.⁵ Compared with the landmark insertion technique, ultrasound guidance showed a higher first-time and overall success rate, lower procedural times, while reducing complications such as arterial puncture, hematomas and pleural puncture.^{3,6-13}

For peripherally inserted central catheters (PICC), ultrasound guidance outperforms also over the landmark technique, with data showing a higher success rate, lower rate of thrombosis and lower costs.¹⁴⁻¹⁸

Ultrasonography-guided peripheral vein cannulation has particularly gained interest among ICU nurses, given that up to one-third of patients have difficult intravenous access resulting from edema, hypovolemia, obesity or previous cannulations.¹⁹ Compared with landmarks, ultrasound guidance has been shown to improve the cannulation rate from 25% to 90%, with fewer attempts, shorter time to access,²⁰⁻²² decreased physician intervention,²³ improved patient satisfaction,²⁴ and reduced the need for a CVC.^{25,26} In veins >12 mm from the skin, using long peripheral venous catheters and midline catheters (MC) improve catheter survival, avoid multiple peripheral vein cannulations and also reduce the need for a CVC.²⁷⁻²⁹ Research has shown that MC may perform similarly to CVC, with a lower rate of catheter-related bloodstream infections.³⁰⁻³²

For arterial cannulation, ultrasound-guided radial artery cannulation showed an increased first-time success rate, overall success rate and lower failure rate, with no differences in the time spent for cannulation or risk of hematoma compared with palpation alone.³³ Ultrasound-guided femoral artery catheterization achieved a 49% reduction in overall complications, including hematoma and accidental venipuncture compared to the landmark technique; it was also associated with a 42% improvement in the likelihood of first-attempt success.³⁴ Other options to radial or femoral artery cannulation, although with less supporting body of evidence, include ultrasound-guided cannulation of the ulnar, axillary, brachial, posterior tibial or dorsalis pedis artery.³⁵⁻³⁸

An intraosseous access is typically indicated in cardiac arrest patients when obtaining an intravenous line is unsuccessful or nonfeasible.³⁹ Options include the proximal tibia, distal tibia, proximal humerus (greater tuberosity), and sternum. Although the needle is often inserted after identification by palpation of the tibial or sternal bones, recognizing the greater tuberosity of the humerus may be challenging. For this indication, ultrasound has demonstrated to be reliable and accurate.⁴⁰

Preprocedure

A high-frequency linear probe is mandatory for vascular cannulation, although a convex probe may be needed in patients with morbid obesity, particularly for femoral vessel targets.⁴¹ A venous or arterial preset is chosen based on

the target vessel, although the preset often requires manual adjustments on a case-by-case basis.

The vessels should be evaluated along both short and long axes. The target vessel should be selected based on patency (e.g., absence of thrombus, complete compressibility and phasic flow in veins; pulsatility, absence of plaques/stenosis and pulsatile flow in arteries), best depth (measuring the distance from the anterior wall of the vessel to the skin, in short axis) and size (measuring the anteroposterior vessel diameter) (Fig. 1A), course (straight better than tortuous) and the safest needle path assessing the relationship of the target vessel with pivotal structures such as the vein, artery or the pleura.⁴² As a rule of thumb, when selecting the catheter size, the anteroposterior diameter of the vessel corresponds to the maximal catheter size (e.g., 5 mm corresponds to a catheter up to 5F). In addition, at least 50% of the catheter length should lie in the vessel; therefore, longer catheters are needed for deeper vessels, and vice versa.

For internal jugular or subclavian vein cannulation, the Trendelenburg position is recommended to enlarge the target veins and minimize the risk of air embolism. However, it may not be safe in some patients, such as those with increased intracranial pressure. In such cases, ultrasound-guided cannulation of the internal jugular vein at a 30° head-up position was demonstrated to be feasible and safe.⁴³ Bilateral lung sliding should also be assessed. To cannulate the femoral vein, the frog-legged position with reverse Trendelenburg aids in increasing the femoral vein size.

Palmar arch sufficiency should be assessed when radial artery cannulation is considered.⁴²

For intraosseous cannulation, as stated previously, the site of needle insertion is often identified by palpation, such as the proximal and distal tibia. However, for insertion into the humerus, ultrasound has demonstrated better performance than palpation in identifying the target at the greater tuberosity.⁴⁰

Procedure

Real-time cannulation technique is preferred. For simplicity, short-axis cannulation is synonymous with the out-of-plane technique, whereas long-axis cannulation is analogous to the in-plane technique.⁴¹ However, practitioners should bear in mind that the out-of-plane and the in-plane terms denote the relationship between the needle and the ultrasound beam, while the short axis and long axis denote the relationship between the ultrasound beam and the vessel. Therefore, it is possible to perform an out-of-plane cannulation in the long-axis or an in-plane cannulation in the short-axis. Either approach is selected on a case-by-case basis.

While a fully sterile technique may be omitted for peripheral venous cannulations, it is mandatory for MC, PICC, arterial, and CVC placements.⁴⁴

After disinfecting the skin and applying a local anesthetic under real-time ultrasound guidance, the needle is inserted in-plane or out-of-plane (Fig. 1B and C), and the catheter is accommodated using the trocar or Seldinger technique.

When using the latter, the guidewire should be confirmed in the vessel before progressing the dilator (Fig. 1D).

To assess the CVC tip position, one operator injects agitated saline or normal saline through the distal port of the CVC, while a second operator performs simultaneously a subcostal 4 chamber view or an apical 4 chamber view. The immediate (within 2 s) appearance of turbulent flow in the right atrium is known as the rapid atrial swirl sign (RASS) and predicts with excellent sensitivity and specificity for a correct catheter tip positioning⁴⁵ (Video 1).

The intravenous position of a peripheral catheter is assessed by direct visualization in the short and/or long axis (Fig. 1E and F) and the flush test (Video 2).

For MC, the catheter is advanced or retroceded until the tip is observed lying in the thoracic tract of the axillary vein.

For arterial cannulation, transient ulnar artery compression may enlarge the radial artery and makes the cannulation easier.⁴⁶ The intra-arterial position of the catheter is assessed by direct visualization in the short and/or long axis.

Intraosseous cannulation is not performed under real-time ultrasound guidance; however, its correct position may be evaluated using color Doppler when subperiosteal flow is observed after flushing the line (Fig. 1G).⁴⁷

Posprocedure

POCUS is performed to assess complications, such as hematoma, pneumothorax (rechecking the lung sliding), infiltration, and catheter-related thrombosis. The distal flow should also be assessed after arterial cannulations, while a pseudoaneurysm should be ruled out (Fig. 1H), when the catheter is removed.

In patients in whom RASS is confirmed and lung sliding is normal, chest radiography can be safely omitted, reducing time, costs, and ionizing radiation exposure.⁴⁸

Table 2 summarizes the steps involved in ultrasound-guided vascular cannulation.

The technique and corresponding ultrasound images of the most commonly cannulated vessels/intraosseous access are provided in the **Supplementary Material, Appendix C**.

Pericardiocentesis

Pericardiocentesis is undoubtedly indicated in tamponade. In patients without hemodynamic compromise, it is indicated for symptomatic moderate-to-large effusion non-responsive to medical therapy, when tuberculous, bacterial, or neoplastic pericarditis is suspected, or in cases of chronic (lasting more than 3 months) large pericardial effusions (>20 mm on echocardiography in diastole).⁴⁹

Blind pericardiocentesis carries morbidity and mortality risks of 20% and 6%, respectively. In contrast, ultrasound-guided pericardiocentesis has demonstrated a success rate of 97%, with a very low rate of complications, and is therefore recommended before, during, and after pericardiocentesis.⁴⁹




Figure 1 Ultrasound-guided vascular cannulation. A) Preprocedural assessment of a peripheral vein for ultrasound-guided cannulation. The anteroposterior diameter of the vein (continuous yellow line, Distancia 2) and distance from the vein to the skin (continuous green line, Distancia 1) are measured. B) In-plane cannulation of the internal jugular vein (v); arrows, needle shaft; arrowhead, needle tip. C) Out-of-plane cannulation of the internal jugular vein (v); a, common carotid artery; arrow, needle. D) The guidewire is observed in the vein lumen (arrows) before placement of a long catheter in the basilic vein (e.g., midline catheter); v, vein. E) A peripheral venous catheter (arrow) is observed in the lumen of a deep vein of the arm (short axis); v, brachial veins; a, brachial artery. F) A peripheral catheter (arrows) is depicted in the lumen of a superficial vein in the long axis; v, vein. G) Subperiosteal flow is observed on color Doppler after intraosseous needle insertion. H) Pseudoaneurysm as a complication of femoral arterial catheterization; black arrows, pseudoaneurysm cavity; white arrows, pseudoaneurysm neck; a, common femoral artery; v, common femoral vein.

Preprocedure

A convex or a phased-array probe is used to assess the view where the effusion accumulation is maximal and closest to the transducer. The thickness of the effusion is measured in diastole in each window, while the distance from the skin to the parietal pericardium and to the myocardium (visceral pericardium) should be obtained to estimate the needle depth of insertion and needle length (Fig. 2A–C). Qualitative assessment should also be performed. While anechoic fluid can be either a transudate or an exudate, the presence of debris or septations points towards the latter. Quantitative and qualitative assessments aid in decision making regarding whether pericardiocentesis, another procedure, or no procedure should be performed. Using a linear probe, the pleura, internal thoracic vessels, and intercostal vessels are delineated to exclude these structures from the needle trajectory (Fig. 2D, E).⁵⁰

Procedure

There are two broad approaches to perform ultrasound-guided pericardiocentesis, which is a real-time procedure: intercostal (apical and left parasternal) and subcostal. The former is the preferred route, given that there is often a shorter distance for the needle to reach the pericardial cavity compared with the subcostal approach. However, it largely depends on the site of maximal fluid accumulation, which is usually non-uniform.⁴⁹

A phased-array probe or convex probe is chosen to guide the procedure. Wherever possible, a linear probe aids in the best observation of the needle, particularly in inter-

costal pericardiocentesis (Fig. 2F). The entire intervention is performed under full sterile barrier precautions.

Pericardiocentesis is performed using a specific kit, or, eventually, in austere settings, a central venous cannulation kit.

After disinfecting and infiltrating the target site with local anesthetics, the needle is inserted in-plane into the pericardial cavity; after aspiration a few milliliters of pericardial fluid, 5 ml of agitated saline are injected through the needle. The immediate appearance of echo contrast in the pericardial space confirms that the needle is in the pericardial cavity and rules out a cardiac chamber perforation⁵¹ (Video 3). The guidewire is then passed and observed in the pericardial space, and, after dilation, the catheter is finally accommodated (Fig. 2G) and connected to a three-way stopcock and a drainage system.

Posprocedure

Rechecking lung sliding is mandatory when using an intercostal approach. If catheter drainage function is optimal, serial ultrasound assessments will show a reduction in the amount of pericardial fluid. When using the subcostal approach, the liver should be observed for a hematoma, and the presence of free peritoneal fluid should also be assessed.

Table 3 summarizes the steps involved in ultrasound-guided pericardiocentesis.

Thoracentesis

Pleural effusions are common in critically ill patients. Although most effusions are of limited clinical significance,

Table 2 Steps for performing ultrasound-guided vascular cannulation.^{8,13,19,27–32,41,42,45}

		Peripheral venous access*	Central venous access	Arterial access
Preprocedure	Patency	- Complete compressibility - Absence of thrombus (anechoic lumen) - Phasic flow on color and spectral Doppler	- Complete compressibility - Absence of thrombus (anechoic lumen) - Phasic flow on color and spectral Doppler	- Anechoic lumen - Pulsatility - Arterial wall without plaques - Pulsatile flow on color and spectral Doppler
	Size	≥4 mm	≥7 mm	Non-reported
	Depth	<12 mm ≥12 mm: consider MC or PICC	As superficial as possible	As superficial as possible
	Other	Delineate surrounding structures (e.g., artery)	Delineate surrounding structures (e.g., artery, pleura) Check bilateral lung sliding in internal jugular vein and/or axillary-subclavian vein cannulations.	Delineate surrounding structures (e.g., vein).
Procedure	Technique	Full sterile barrier technique. Skin disinfection and local anesthesia infiltration before puncture. An out-of-plane or in-plane real-time approach is selected on a case- by -case basis.		
	Catheter “tips”	Observe the catheter into the vessel and use the flush test. For MC, assess catheter tip position in the thoracic tract of the axillary vein. Each time the Seldinger technique is used, the guidewire should be observed into the vessel prior to passing the dilator.	Each time the Seldinger technique is used, the guidewire should be observed into the vessel prior to passing the dilator. Assess catheter tip position with the rapid atrial swirl sign (RASS) using 10 ml of normal or agitated saline.	Observe the catheter into the artery lumen.
Posprocedure		Assess for thrombosis, inadvertent arterial cannulation, hematomas, infiltration.	Assess for thrombosis, hematomas, inadvertent arterial cannulation, infiltration. Recheck lung sliding. If bilateral lung sliding is normal and RASS is observed, chest radiograph can be omitted.	Assess for thrombosis, inadvertent venous cannulation, hematomas, pseudoaneurysm, assess distal flow.

* Superficial or epifascial veins (e.g., basilic, cephalic) are distinguished from deep veins (e.g., brachial veins) given that the former does not run along with artery or nerve. RASS, rapid atrial swirl sign; MC, midline catheter; PICC, peripherally inserted central catheter.

aggressive management is required in some cases. These are broadly classified as transudates or exudates. The causes of transudative effusions include volume overload, hypoalbuminemia, and regions of altered pleural pressure attributable to atelectasis and mechanical ventilation. Exudates may be caused by pulmonary or pleural infections, pulmonary embolism, postsurgical complications, and malignancy. Increases in pleural fluid volume are accommodated principally by chest wall expansion and, to a lesser degree, by lung collapse.⁵² When indicated, diagnostic thoracocen-

tesis is key to distinguishing transudative effusions from exudative effusions. In addition, when large, pleural effusions may compromise respiratory function and patients may benefit from drainage.

A large body of evidence shows that when compared with the landmark technique, ultrasound-guided thoracentesis improves the success rate, decreases costs, reduces the rate of pneumothorax by 19%, and minimizes the occurrence of hemothorax.^{53,54}




Figure 2 Ultrasound-guided pericardiocentesis. A) Subcostal 4 chamber view. B) Parasternal long-axis view. C) Apical view. As noted, the optimal window to insert the needle is the apical, given the shorter distance to reach the pericardial space (continuous green line) and higher pericardial fluid thickness (continuous yellow line). RA, right atrium; RV, right ventricle; LA, left atrium; LV, left ventricle; RVOT, right ventricular outflow tract. Asterisks indicate pericardial effusion. Adapted from Blanco P, Figueroa L, Menéndez MF, Berrueta B. Pericardiocentesis: ultrasound guidance is essential. *Ultrasound J.* 2022;14(1):9. <https://theultrasoundjournal.springeropen.com/articles/10.1186/s13089-022-00259-5>. (CC-BY-4.0). D) Recognition of the left internal thoracic vessels along the left parasternal line (dotted white line) with a linear probe on two-dimensional and color Doppler imaging; s-sct, skin-subcutaneous tissue; m, intercostal muscle; r, rib; arrowhead, pleura; asterisks, internal thoracic vessels. E) Recognition of the intercostal vessels with a linear probe on color Doppler imaging. Real-time in-plane ultrasound-guided pericardiocentesis via intercostal approach (apical view) using a linear probe. The needle (arrows) is entirely observed in the pericardial space (asterisks); LV: left ventricle. G) Hemorrhagic fluid is freely evacuated from the pericardial space after catheter placement.

Table 3 Steps for performing ultrasound-guided pericardiocentesis.⁴⁹⁻⁵¹

Preprocedure	<ul style="list-style-type: none"> Determine the presence of the pericardial effusion Estimate the quantity of the effusion (thickness in diastole) Measure the distance from the skin to parietal pericardium and from the skin to the myocardium Assess for the presence of septations, debris Delineate the intercostal and internal thoracic vessels Delineate the pleura and assess lung sliding
Procedure	<ul style="list-style-type: none"> Use in-plane real-time technique Thoracic approaches are preferred over subcostal approach Skin disinfection and local anesthesia infiltration before puncture. When using intercostal approaches, the needle should be inserted along the superior border of the rib as a way to prevent inadvertent injury to the intercostal vessels, which runs along the inferior rib border. When the needle is observed in the pericardial space and fluid is freely aspirated, injectate 5 ml of agitated saline and observe the bubbles in the pericardial space Then progress the guidewire, which should be observed in the pericardial space before passing the dilator Pass the catheter and connect to a drainage system
Posprocedure	<ul style="list-style-type: none"> Re-check lung sliding. Re-check the remaining pericardial fluid. When using a subcostal approach, check for liver hematomas and free peritoneal fluid

Preprocedure

The patient should lie supine in a slight semi-recumbent position. With a convex or phased-array probe, it is of paramount relevance to estimate the size of the pleural effusion to define whether thoracentesis is needed or safe, another procedure, or no procedure should be performed. While often eyeballed, a drainable pleural effusion may be defined when the distance between the visceral and parietal pleura is ≥ 10 mm.⁵⁵ In addition, the ultrasonographic characteristics of the effusion should also be assessed based on the presence of septations, swirling debris (i.e., plankton sign) or pleural thickening. While anechoic effusions can be either transudative or exudative, those showing one or more of the aforementioned signs are often exudative and should prompt pleural fluid sampling (Fig. 3A and B).⁵³ The distance from the skin to the parietal and visceral pleura should be measured using ultrasound to aid in the selection of an adequate needle length and estimate the depth of insertion (Fig. 3C). The diaphragm must also be delineated. Chest wall vessels should be excluded from the needle trajectory, which is assessed with a linear probe using color Doppler (Fig. 3D).⁵⁶ Bilateral lung sliding should also be determined prior to the procedure.

Procedure

The procedure can be performed under static or real-time ultrasound guidance. Although both techniques have proven to be useful, the static technique is preferable. The added effort and complexity of the real-time technique is rarely justified, as moderate or large effusions are easy to access, provided the pre-procedure scan is made correctly.⁵⁶ Real-time guidance is best reserved for small or loculated effusions, where strict control of needle trajectory and depth is mandatory.⁵⁶

With the static technique, and with the patient lying in the same position as the preprocedural scan, the best insertion site is selected with ultrasound and is marked with a pen on the skin (Fig. 3E). After disinfecting and infiltrating the target site with local anesthetics, the needle is advanced until pleural fluid is obtained. The needle should be inserted along the superior border of the rib to prevent inadvertent injury to the intercostal vessels that run along the inferior rib border.

In the dynamic technique, the needle is inserted in-plane and observed in real-time, entering the pleural space when pleural fluid is aspirated (Fig. 3F). A fully sterile technique is required to do so.

When evacuation of the effusion is needed, a pig-tail catheter is typically inserted.⁵⁷ In patients with non-complicated pleural effusions, pleural drainage using central venous catheters has proven useful and safe.⁵⁸ Each time the Seldinger technique is used, the guidewire should be observed in the pleural space before passing the dilator (Fig. 3G). Then, the catheter is advanced and the guidewire is removed. Intrapleural placement of the catheter can also be confirmed by ultrasonography, following its course from the superficial soft tissues to the pleural cavity (Fig. 3H). Finally, the catheter is connected to a drainage system,

which could be a water-seal chest drainage or a urinary bag collection system,⁵⁸ interposing a Heimlich valve.

Posprocedure

Lung ultrasound is performed to rule out pneumothorax. A comparison with preprocedural scanning is helpful. Among the several well known ultrasonographic signs of pneumothorax, the hydro-point sign should also be actively investigated.⁵⁹ The hydro-point defines the presence of hydropneumothorax and is best assessed with a convex or phased-array probe in basal lateral or posterolateral views of the thorax.^{60,61} This sign is depicted as contact between the pleural fluid and the artifacts of the intrapleural air, which does not show lung sliding⁶¹ (Video 4). Chest radiography may be safely omitted in patients in whom a successful thoracentesis is performed and no ultrasonographic signs of pneumothorax are detected.

Table 4 summarizes the steps involved in ultrasound-guided thoracentesis.

Paracentesis

Free peritoneal fluid (i.e., ascites) is frequent in critical care patients, with etiologies ranging from fluid overload, cardiac or hepatic failure, peritonitis, etc. Several times, there is a need for fluid investigation (transudate vs. exudate) and also fluid evacuation to improve respiratory function and reduce intra-abdominal pressure. The most common serious complication of paracentesis is bleeding (abdominal wall hematoma and hemoperitoneum) through laceration of the abdominal wall vessels (e.g., inferior epigastric arteries or its branches), although puncture of the bowel and other abdominal organs has also been observed. Compared with the landmark technique, ultrasound guidance reduced the risk of bleeding by 68%,⁵⁴ while improving the success rate to 95%–100%, as opposed to 65% for the landmark technique.⁶²

Preprocedure

Using a convex or phased-array probe, POCUS aids in detecting the presence, volume, and characteristics of the peritoneal free fluid (anechoic fluid can be either transudative or exudative, while the presence of septations or debris indicates an exudative effusion),⁶³ and determines whether a diagnostic or therapeutic paracentesis is needed or safe, another procedure, or no procedure should be performed.

The maximal site of fluid accumulation is often eyeballed by ultrasonography (although the smallest fluid depth can be used to quantify the effusion), and the distance from the skin to the parietal peritoneum and visceral peritoneum is measured to aid in the selection of the needle length and estimate the depth of insertion (Fig. 4A).⁶³

Using a linear probe, the abdominal wall vessels should also be identified and excluded from the needle trajectory (Fig. 4B).

The patient should lie in a slightly semi-recumbent position.




Figure 3 Ultrasound-guided thoracentesis. A) A simple pleural effusion (asterisks), diaphragm (d) and lung (L) are delineated using ultrasound. B) A complex pleural effusion (asterisk) is observed; d, diaphragm. C) The best fluid pocket for thoracentesis is selected by measuring the distance from the skin to the pleural effusion (asterisk) and the effusion depth; L, lung; d, diaphragm. D) The intercostal vessels are delineated before cannulation using a linear transducer and color Doppler. E) The insertion site is marked on the skin. F) Dynamic ultrasound guidance for thoracentesis. Arrows, needle shaft; arrowhead, needle tip; asterisk, pleural effusion; L, lung. G) The guidewire (arrows) is observed within the pleural effusion (asterisks); d, diaphragm. H) A central catheter (arrows) is observed within the pleural effusion (asterisks); L, lung.

Table 4 Steps for performing ultrasound-guided thoracentesis.^{53,55,56,57-61}

Preprocedure	Determine the presence of the pleural effusion Estimate the quantity of the pleural effusion (often eyeballed) Measure the distance from the skin to the parietal pleura and to the lung Assess for the presence of septations, debris, pleural thickening Delineate the diaphragm Delineate the intercostal vessels Check the lung sliding
Procedure	The static technique is of first choice Skin disinfection and local anesthesia infiltration before puncture. Use dynamic technique in small-sized or loculated effusions When using the Seldinger technique, check whether the guidewire is in the pleural cavity before passing the dilator/catheter Pass the catheter and connect to a drainage system
Posprocedure	Re-check lung sliding and search for other signs of pneumothorax, particularly the hydro-point sign. Re-check the remaining pleural fluid.

Procedure

Either static or real-time technique can be used. Similar to ultrasound-guided thoracentesis, the static technique is preferred. The insertion site should be marked immediately before performing the procedure (Fig. 4C), and the patient should remain in the same position between marking the site and performing the procedure, given that free-flowing peritoneal fluid and abdominal organs, especially loops of the small bowel, can easily shift when a patient changes position or takes a deep breath.⁶²

Real-time ultrasound-guided paracentesis (in-plane or out-of-plane) (Fig. 4D) is best suited for obese patients, for patients with small fluid collections, or when performing the procedure near critical structures, such as loops of the small bowel, liver, or spleen.⁶²

When there is a need to evacuate fluid, a pigtail catheter is preferred, although a central catheter can be used in selected cases.⁶⁴ The catheter position could also be confirmed using ultrasound (Fig. 4E).

Posprocedure

The amount of remaining peritoneal fluid should be assessed. The presence of an abdominal wall hematoma can also be delineated using ultrasound.

Table 5 summarizes the steps involved in ultrasound-guided paracentesis.




Figure 4 Ultrasound-guided paracentesis. A) Ascites is observed (asterisk), and the distance from the skin to the effusion and to a bowel loop (b) is obtained. B) The inferior epigastric vessels (boxes) are delineated into the abdominis rectus sheath using a linear probe on two-dimensional and color Doppler imaging; s, skin; sct, subcutaneous tissue; m, rectus abdominis muscle; asterisk, ascites. C) The insertion site is marked on the skin, and puncture is performed under static guidance. D) Real-time ultrasound-guided paracentesis; arrows, needle shaft; arrowhead, needle tip; asterisk, ascites. E) A locking pigtail catheter is observed within the ascites (asterisks). The arrow indicates the body, whereas the arrowhead indicates the tip of the pigtail.

Table 5 Steps for performing ultrasound-guided paracentesis.^{62,63}

Preprocedure	Determine the presence of free peritoneal fluid Estimate the quantity of free peritoneal fluid Measure the distance from the skin to the parietal peritoneum and to the bowel Assess for the presence of septations, debris Delineate the diaphragm
Procedure	Delineate the vessels of the abdominal wall The static technique is of first choice Use dynamic techniques in small-sized or loculated effusions Skin disinfection and local anesthesia infiltration before puncture When using the Seldinger technique, check whether the guidewire is in the peritoneal cavity before passing the dilator/catheter Pass the catheter and connect to a drainage system
Posprocedure	Re-check the remaining free-peritoneal fluid Assess for abdominal wall hematomas

Soft-tissue fluid collection aspiration/arthrocentesis

Ultrasound has demonstrated high accuracy in differentiating superficial abscesses from cellulitis, with a sensitivity of 96.2%, specificity of 82.9%, positive likelihood ratio of 5.63, and negative likelihood ratio of 0.05.⁶⁵ Cellulitis is observed as a cobblestoning pattern of the subcutaneous tissue, whereas an abscess is a heterogeneous fluid

collection.⁶⁶ Differentiating an abscess from other fluid collections (e.g., hematoma) is challenging; ultrasound-guided diagnostic needle aspiration may be useful in such cases.⁶⁶ Furthermore, aspiration of small abscesses (<4 cm) may avoid incision and drainage (I&D) in some cases.⁶⁷ However, aspiration of abscess with septations or satellite lesions may not be feasible despite the use of ultrasound guidance, as reported for Methicillin-resistant *Staphylococcus aureus* infections.⁶⁸ In other cases, ultrasound aids in the

best definition of the site for I&D compared with physical examination alone.⁶⁹

Arthrocentesis is mainly indicated when septic arthritis (SA) is highly suspected. The incidence of SA is 4–21 per 100,000 person/years.⁷⁰ Large and small joints can be involved; however, the knee is most frequently affected (50% of cases). Monoarticular involvement is the most common presentation; however, up to 20% of cases may have concurrent infection in several joints.⁷⁰ Ultrasound-guided knee arthrocentesis provides greater accuracy and clinical improvement than the landmark technique.⁷¹ Other joint effusions can also be best delineated and aspirated with ultrasound guidance when compared with landmarks such as the hip, elbow, wrist, or ankles.^{72–74}

Preprocedure

With a linear probe (or a convex probe in obese or edematous patients), ultrasound aids in differentiating subcutaneous tissue edema, soft-tissue fluid collections (e.g., abscesses), periarticular disease (e.g., bursitis), or an intra-articular effusion.

In transverse and longitudinal views, soft-tissue fluid collections are measured, as well as the distance to the skin, to estimate the needle depth of insertion and to choose an adequate needle length (Fig. 5A,B). Color Doppler aids in defining the presence of vascular structures near the lesion, to avoid puncturing when inserting the needle.

Ultrasound anatomy of each joint should be well known for assessing effusion. Given its frequency and ease of access, knee ultrasound is reviewed herein. With the linear probe placed longitudinally above the patella and the probe marker pointing cephalad, the superior portion of the patella, quadriceps femoris tendon (long axis), fat pad, and femur should be observed (Fig. 5C). The effusion will be located below the quadriceps tendon and above the femur (Fig. 5D). Once fluid effusion is recognized, the probe is rotated 90 degrees into a transverse view so that the probe marker faces to the patient's right side. The distance from the skin to the effusion is then measured to estimate the needle depth of insertion and length to be used.

Procedure

Real-time technique is used to aspirate soft-tissue collections and perform arthrocentesis. The out-of-plane and in-plane approaches are selected on a case-by-case basis. For example, aspiration of abscess and knee arthrocentesis are best performed using the in-plane approach (Fig. 5E and 5F), whereas the out-of-plane approach is preferred for aspiration of effusion from the wrist joint. A fully sterile barrier technique is required to do so.

Posprocedure

Investigation of post-procedural soft-tissue hematomas should be ruled in or out, as well as reassessing the joint effusion size and the presence of hemarthrosis, comparing with pre-procedural evaluation.

The technique for ultrasound-guided arthrocentesis is provided in the **Supplementary Material, Appendix D**.

Lumbar puncture

Lumbar puncture (LP) is often performed in critical care patients to rule in or out meningitis, Guillain-Barre syndrome, and in selected cases for providing spinal or epidural anesthesia. Although many practitioners successfully perform the procedure using anatomical landmarks, the failure rate reaches up to 19%.⁷⁵ Ultrasound guidance has demonstrated to significantly reduce the risk of traumatic taps, number of insertion attempts, and number of needle redirections,⁷⁶ with the greatest benefits demonstrated in patients with poorly palpable landmarks, as in obesity, and in patients with scoliosis, previous spinal surgery, or a history of difficult LP.⁷⁷

Preprocedure

With the sitting patient or in a lateral decubitus (the latter often in critical care patients), a convex probe is placed in the transverse plane (perpendicular to the long axis of the spine) in the midline of the lower back. Here, the spinal processes (SPs) are recognized as superficial hyperechoic peaked structures with posterior acoustic shadowing (Fig. 6A), while the laminae or articular processes ("bat sign") are seen as hyperechoic structures deep and lateral to the SPs. The hypoechoic interspinal spaces can be delineated by moving the transducer cephalad or caudally to the SPs. With the SPs centered on the ultrasound screen, the skin is marked with a pen at the transducer midpoint over several vertebrae (Fig. 6A).

The transducer is then placed longitudinally (parallel to the long axis of the spine). Again, the SPs are observed as bright peaked structures with posterior acoustic shadowing; between them, there are the hypoechoic interspinal spaces (Fig. 6B). To recognize the sacrum, the transducer is moved caudally until a roughly hyperechoic line is observed. With the interspinal space centered on the ultrasound screen, the skin is marked with a pen at the transducer midpoint over several vertebrae.

After marking the skin in the transverse and longitudinal planes, both lines are continued until they intersect the midline, where the puncture will be performed over the broad spinal space (Fig. 6C).

Slightly angling the transducer to the midline, the paramedian longitudinal approach is obtained (Fig. 6D). Here, the pedicles and posterior complex (including the ligamentum flavum, epidural space, and posterior dura) are easily recognized (Fig. 6D) Measuring the distance from the skin to the posterior complex aids in estimating the needle depth of insertion and length to be used.(Fig. 6D)

Procedure

The procedure can be performed with the patient sitting (preferably) or lying in a lateral decubitus position.




Figure 5 Ultrasound-guided aspiration of soft tissue collections and arthrocentesis. A) Soft tissue fluid collection is identified and measured. B) Distances are measured to estimate the needle depth of insertion and length to be used. C) Technique for finding the suprapatellar articular recess of the knee and its corresponding ultrasound images; p, patella; t, cuadriceps tendon; f, femur; asterisk, fat pad (articular recess). D) Knee effusion is indicated by the symbol #. E) Real-time ultrasound-guided arthrocentesis of the knee, obtaining a purulent fluid. The arrowhead indicates the level of the needle that reached the articular effusion. F) Real-time ultrasound-guided aspiration of soft tissue collection. The needle is indicated by arrows.

The use of the static technique is largely supported in the literature; however, real-time technique have shown comparable or even better outcomes.⁷⁸

The static procedure technique is performed following the mark on the skin, advancing the needle until a loss of resistance is felt and the cerebrospinal fluid (CSF) starts coming out.

For the real-time technique, the LP is performed using the paramedian approach. The needle is inserted in-plane, until it reaches and passes through the posterior complex when a loss of resistance is felt, and CSF fluid is obtained.




Figure 6 Ultrasound-guided lumbar puncture. A) With the linear transducer placed transversally over the midline of the lower back, the spinal processes (sp) are recognized and the skin marked with a pen. B) After rotating the transducer 90 degrees clockwise and placing it longitudinally over the midline of the lower back, the spinal processes (sp) are seen, and the interspinous space (arrow) is delineated, and the skin is marked with a pen. C) Intersecting the skin markings obtained in A) and B), the needle insertion site is marked with a pen. D) Parasagittal approach and the corresponding ultrasound image. To obtain this view, the transducer is slightly basculated to one side from the longitudinal midline view. In the ultrasound image, the spinal muscles (spm), laminae (L), posterior complex (white arrows), and posterior longitudinal ligament (black arrows) are well recognized. The distance to the posterior complex is measured (dotted green line), to approximate the needle depth of insertion during the lumbar puncture.

Posprocedure

Post-procedural evaluation is mainly clinical, such as the presence of a visible hematoma on the skin, postdural puncture headache, back pain, and eventually spinal hematoma, which is rare but devastating complication of LP. While there are publications in pediatrics for recognizing epidural hematomas using ultrasound,⁷⁹ there are no data on the adult population in whom magnetic resonance imaging is typically indicated.

Table 6 summarizes the steps involved in ultrasound-guided lumbar puncture.

Future directions

Newer ultrasound technologies may be applied for procedural guidance, such as biplane imaging (also known as Xplane®), which simultaneously shows two-orthogonal

planes. Matrix array transducers are required for this purpose, although they are typically not available in the ultrasound machines of the ICU. Anecdotal evidence using a miniaturized device with biplane imaging has shown its usefulness for vascular cannulation.⁸⁰

Conclusions

Ultrasound guidance is key to guaranteeing the success and safety of several procedures performed in the ICU. With time, the procedures performed by intensivists may extend beyond the core competencies depicted in this review.

Funding

This work has not been supported by any grants.

Table 6 Steps for performing ultrasound-guided lumbar puncture.^{75,76}

Preprocedure	Use a linear or a convex probe Place the transducer in the midline in the longitudinal and transverse planes and mark on the skin over several vertebrae. Continues the lines until intersect in the midline Measure the distance from the skin to the flavum ligament on a longitudinal paramedian approach to estimate the needle depth of insertion and needle length
Procedure	The static technique is of first choice Real-time technique from a longitudinal paramedian approach can also be performed Skin disinfection and local anesthesia infiltration before puncture When the posterior complex is passed with the needle, cerebrospinal fluid is obtained.
Posprocedure	Clinical evaluation is mainly performed: skin hematomas, postpuncture dural headache, back pain and spinal hematoma (rare) for which cerebral magnetic resonance imaging is mandatory.

Conflicts of interest

Authors have no conflicts of interest to disclose.

This work has not been presented at any conferences.

Authors' contributions

Anselmo Abdo-Cuza (AAC) revised the manuscript for important intellectual content and provided raw material for creating the figures and editing the videos. AAC read and approved the final manuscript.

Cristina Martínez Díaz (CMD) revised the manuscript for important intellectual content and provided raw material for creating the figures and editing the videos. CMD read and approved the final manuscript.

Elena Abril Palomares (EAP) revised the manuscript for important intellectual content and provided raw material for creating the figures and editing the videos. EAP read and approved the final manuscript.

Pablo Blanco (PB) performed the literature review, wrote the drafts, coordinated the input of all authors, created figures/edited the videos, and revised the manuscript and appendices for important intellectual content. PB read and approved the final manuscript.

Virginia Fraile Gutiérrez (VFG) revised the manuscript for important intellectual content and provided raw material for creating the figures and editing the videos. VFG read and approved the final manuscript.

Appendix A Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.medint.2023.05.008>.

References

- Blanco P, Miralles Aguiar F, Vallejo A. Point-of-care ultrasonography in critical care medicine: a one way directional road. *J Ultrasound*. 2016;19(2):157–8, <http://dx.doi.org/10.1007/s40477-015-0192-z>.
- Robba C, Wong A, Poole D, Al Tayar A, Arntfield RT, Chew MS, et al. Basic ultrasound head-to-toe skills for intensivists in the general and neuro intensive care unit population: consensus and expert recommendations of the European Society of Intensive Care Medicine. *Intensive Care Med*. 2021;47(12):1347–67, <http://dx.doi.org/10.1007/s00134-021-06486-z>.
- Nicolaou S, Talsky A, Khashoggi K, Venu V. Ultrasound-guided interventional radiology in critical care. *Crit Care Med*. 2007;35 5 Suppl:5186–197, <http://dx.doi.org/10.1097/01.ccm.0000260630.68855.df>.
- Barr L, Hatch N, Roque PJ, Wu TS. Basic ultrasound-guided procedures. *Crit Care Clin*. 2014;30(2):275–304, <http://dx.doi.org/10.1016/j.ccc.2013.10.004>, vi.
- Zieleskiewicz L, Muller L, Lakhil K, Meresse Z, Arbelot C, Bertrand PM, et al. Point-of-care ultrasound in intensive care units: assessment of 1073 procedures in a multicentric, prospective, observational study. *Intensive Care Med*. 2015;41(9):1638–47, <http://dx.doi.org/10.1007/s00134-015-3952-5>.
- Hind D, Calvert N, McWilliams R, Davidson A, Paisley S, Beverley C, et al. Ultrasonic locating devices for central venous cannulation: meta-analysis. *BMJ*. 2003;327(7411):361, <http://dx.doi.org/10.1136/bmj.327.7411.361>.
- Kwon TH, Kim YL, Cho DK. Ultrasound-guided cannulation of the femoral vein for acute haemodialysis access. *Nephrol Dial Transplant*. 1997;12(5):1009–12, <http://dx.doi.org/10.1093/ndt/12.5.1009>.
- Bouaziz H, Zetlaoui PJ, Pierre S, Desruennes E, Fritsch N, Jochum D, et al. Guidelines on the use of ultrasound guidance for vascular access. *Anaesth Crit Care Pain Med*. 2015;34(1):65–9, <http://dx.doi.org/10.1016/j.accpm.2015.01.004>.
- Brass P, Hellmich M, Kolodziej L, Schick G, Smith AF. Ultrasound guidance versus anatomical landmarks for subclavian or femoral vein catheterization. *Cochrane Database Syst Rev*. 2015;1(1):CD011447, <http://dx.doi.org/10.1002/14651858.cd011447>.
- Lalu MM, Fayad A, Ahmed O, Bryson GL, Fergusson DA, Barron CC, et al. Ultrasound-guided subclavian vein catheterization: a systematic review and meta-analysis. *Crit Care Med*. 2015;43(7):1498–507, <http://dx.doi.org/10.1097/ccm.0000000000000973>.
- O'Leary R, Ahmed SM, McLure H, Oram J, Mallick A, Bhambra B, et al. Ultrasound-guided infraclavicular axillary vein cannulation: a useful alternative to the internal jugular vein. *Br J Anaesth*. 2012;109(5):762–8, <http://dx.doi.org/10.1093/bja/aes262>.
- Gualtieri E, Deppe SA, Sippertly ME, Thompson DR. Subclavian venous catheterization: greater success rate for less experienced operators using ultrasound guidance. *Crit Care Med*. 1995;23(4):692–7, <http://dx.doi.org/10.1097/00003246-199504000-00018>.

13. Lamperti M, Biasucci DG, Disma N, Pittiruti M, Breschan C, Vailati D, et al. European Society of Anaesthesiology guidelines on peri-operative use of ultrasound-guided for vascular access (PERSEUS vascular access). *Eur J Anaesthesiol.* 2020;37(5):344–76, <http://dx.doi.org/10.1097/eja.0000000000001180>.
14. Sofocleous CT, Schur I, Cooper SG, Quintas JC, Brody L, Shelin R. Sonographically guided placement of peripherally inserted central venous catheters: review of 355 procedures. *AJR Am J Roentgenol.* 1998;170(6):1613–6, <http://dx.doi.org/10.2214/ajr.170.6.9609183>.
15. Nichols I, Humphrey JP. The efficacy of upper arm placement of peripherally inserted central catheters using bedside ultrasound and microintroducer technique. *J Infus Nurs.* 2008;31(3):165–76, <http://dx.doi.org/10.1097/01.nan.0000317703.66395.b8>.
16. Stokowski G, Steele D, Wilson D. The use of ultrasound to improve practice and reduce complication rates in peripherally inserted central catheter insertions: final report of investigation. *J Infus Nurs.* 2009;32(3):145–55, <http://dx.doi.org/10.1097/nan.0b013e3181a1a98f>.
17. Robinson MK, Mogensen KM, Grudinskas GF, Kohler S, Jacobs DO. Improved care and reduced costs for patients requiring peripherally inserted central catheters: the role of bedside ultrasound and a dedicated team. *JPEN J Parenter Enteral Nutr.* 2005;29(5):374–9, <http://dx.doi.org/10.1177/0148607105029005374>.
18. Li J, Fan YY, Xin MZ, Yan J, Hu W, Huang WH, et al. A randomised, controlled trial comparing the long-term effects of peripherally inserted central catheter placement in chemotherapy patients using B-mode ultrasound with modified Seldinger technique versus blind puncture. *Eur J Oncol Nurs.* 2014;18(1):9, <http://dx.doi.org/10.1016/j.ejon.2013.08.003>.
19. Blanco P. Ultrasound-guided peripheral venous cannulation in critically ill patients: a practical guideline. *Ultrasound J.* 2019;11(1):27, <http://dx.doi.org/10.1186/s13089-019-0144-5>.
20. Costantino TG, Parikh AK, Satz WA, Fojtik JP. Ultrasonography-guided peripheral intravenous access versus traditional approaches in patients with difficult intravenous access. *Ann Emerg Med.* 2005;46(5):456–61, <http://dx.doi.org/10.1016/j.annemergmed.2004.12.026>.
21. Gregg SC, Murthi SB, Sisley AC, Stein DM, Scalea TM. Ultrasound-guided peripheral intravenous access in the intensive care unit. *J Crit Care.* 2010;25(3):514–9, <http://dx.doi.org/10.1016/j.jcrc.2009.09.003>.
22. Mahler SA, Wang H, Lester C, Conrad SA. Ultrasound-guided peripheral intravenous access in the emergency department using a modified Seldinger technique. *J Emerg Med.* 2010;39(3):325–9, <http://dx.doi.org/10.1016/j.jemermed.2009.02.013>.
23. Bauman M, Braude D, Crandall C. Ultrasound-guidance vs. standard technique in difficult vascular access patients by ED technicians. *Am J Emerg Med.* 2009;27(2):135–40, <http://dx.doi.org/10.1016/j.ajem.2008.02.005>.
24. Schoenfeld E, Shokoohi H, Boniface K. Ultrasound-guided peripheral intravenous access in the emergency department: patient-centered survey. *West J Emerg Med.* 2011;12(4):475–7, <http://dx.doi.org/10.5811/westjem.2011.3.1920>.
25. Au AK, Rotte MJ, Grzybowski RJ, Ku BS, Fields JM. Decrease in central venous catheter placement due to use of ultrasound guidance for peripheral intravenous catheters. *Am J Emerg Med.* 2012;30(9):1950–4, <http://dx.doi.org/10.1016/j.ajem.2012.04.016>.
26. Galen B, Baron S, Young S, Hall A, Berger-Spivack L, Southern W. Reducing peripherally inserted central catheters and midline catheters by training nurses in ultrasound-guided peripheral intravenous catheter placement. *BMJ Qual Saf.* 2020;29(3):245–9, <http://dx.doi.org/10.1136/bmjqs-2019-009923>.
27. Pandurangadu AV, Tucker J, Brackney AR, Bahl A. Ultrasound-guided intravenous catheter survival impacted by amount of catheter residing in the vein. *Emerg Med J.* 2018;35(9):550–5, <http://dx.doi.org/10.1136/emered-2017-206803>.
28. Bahl A, Hang B, Brackney A, Joseph S, Karabon P, Mohammad A, et al. Standard long IV catheters versus extended dwell catheters: a randomized comparison of ultrasound-guided catheter survival. *Am J Emerg Med.* 2019;37(4):715–21, <http://dx.doi.org/10.1016/j.ajem.2018.07.031>.
29. Bahl A, Hijazi M, Chen NW, Lachapelle-Clavette L, Price J. Ultralong versus standard long peripheral intravenous catheters: a randomized controlled trial of ultrasonographically guided catheter survival. *Ann Emerg Med.* 2020;76(2):134–42, <http://dx.doi.org/10.1016/j.annemergmed.2019.11.013>.
30. Prasanna N, Yamane D, Haridasa N, Vasovator D, Sparks A, Hawkins K. Safety and efficacy of vasopressor administration through midline catheters. *J Crit Care.* 2021;61:1–4, <http://dx.doi.org/10.1016/j.jcrc.2020.09.024>.
31. Tripathi S, Kumar S, Kaushik S. The practice and complications of midline catheters: a systematic review. *Crit Care Med.* 2021;49(2):e140–50, <http://dx.doi.org/10.1097/ccm.0000000000004764>.
32. Blanco P, Figueroa L, Menéndez MF, Berrueta B. The midline venous catheter in critically ill COVID-19 patients. *Med Intensiva.* 2022;46(10):591–3, <http://dx.doi.org/10.1016/j.medine.2021.09.002>.
33. Moussa Pacha H, Alahdab F, Al-Khadra Y, Idris A, Rabbat F, Dar-moch F, et al. Ultrasound-guided versus palpation-guided radial artery catheterization in adult population: a systematic review and meta-analysis of randomized controlled trials. *Am Heart J.* 2018;204:1–8, <http://dx.doi.org/10.1016/j.ahj.2018.06.007>.
34. Sobolev M, Slovut DP, Lee Chang A, Shiloh AL, Eisen LA. Ultrasound-guided catheterization of the femoral artery: a systematic review and meta-analysis of randomized controlled trials. *J Invasive Cardiol.* 2015;27(7):318–23. Retrieved December 15, 2022 from <https://www.hmpgloballearningnetwork.com/site/jic/articles/let-us-guide-way>
35. Vidovich MI. Ulnar artery catheterization: is this our second access site or is it still femoral? *Curr Cardiol Rep.* 2018;20(10):91, <http://dx.doi.org/10.1007/s11886-018-1039-y>.
36. Flumignan RL, Trevisani VF, Lopes RD, Baptista-Silva JC, Flumignan CD, Nakano LC. Ultrasound guidance for arterial (other than femoral) catheterisation in adults. *Cochrane Database Syst Rev.* 2021;10(10):CD013585, <http://dx.doi.org/10.1002/14651858.cd013585.pub2>.
37. Htet N, Vaughn J, Adigopula S, Hennessey E, Mihm F. Needle-guided ultrasound technique for axillary artery catheter placement in critically ill patients: a case series and technique description. *J Crit Care.* 2017;41:194–7, <http://dx.doi.org/10.1016/j.jcrc.2017.05.026>.
38. Anand RK, Maitra S, Ray BR, Baidhya DK, Khanna P, Chowdhury SR, et al. Comparison of ultrasound-guided versus conventional palpation method of dorsalis pedis artery cannulation: a randomized controlled trial. *Saudi J Anaesth.* 2019;13(4):29, http://dx.doi.org/10.4103/sja.sja_766_18.
39. Panchal AR, Bartos JA, Cabañas JG, Donnino MW, Drennan IR, Hirsch KG, et al. Part 3: adult basic and advanced life support: 2020 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation.* 2020;142 16_suppl_2:S366–468, <http://dx.doi.org/10.1161/cir.0000000000000916>.
40. Bustamante S, Bajracharya GR, Cheruku S, Leung S, Mao G, Singh A, et al. Point-of-care ultrasound to identify landmarks of the proximal humerus: potential use for intraosseous

- vascular access. *J Ultrasound Med.* 2021;40(4):725–30, <http://dx.doi.org/10.1002/jum.15442>.
41. Franco-Sadud R, Schnobrich D, Mathews BK, Candotti C, Abdel-Ghani S, Perez MG, et al. Recommendations on the use of ultrasound guidance for central and peripheral vascular access in adults: a position statement of the Society of Hospital Medicine. *J Hosp Med.* 2019;14:E1–22, <http://dx.doi.org/10.12788/jhm.3287>.
 42. Blanco P. Ultrasound-guided vascular cannulation in critical care patients: a practical review. *Blanco. Med Intensiva.* 2016;40(9):560–71, <http://dx.doi.org/10.1016/j.medin.2016.07.009>.
 43. Bredertau J, Greim C, Schwemmer U, Haunschmid B, Markus C, Roewer N. Ultrasound-guided cannulation of the internal jugular vein in critically ill patients positioned in 30 degrees dorsal elevation. *Eur J Anaesthesiol.* 2004;21(9):684–7, <http://dx.doi.org/10.1017/s0265021504009032>.
 44. Millington SJ, Hendin A, Shiloh AL, Koenig S. Better with ultrasound: peripheral intravenous catheter insertion. *Chest.* 2020;157(2):369–75, <http://dx.doi.org/10.1016/j.chest.2019.04.139>.
 45. Korsten P, Mavropoulou E, Wienbeck S, Ellenberger D, Patschan D, Zeisberg M, et al. The “rapid atrial swirl sign” for assessing central venous catheters: performance by medical residents after limited training. *PLoS One.* 2018;13(7):e0199345, <http://dx.doi.org/10.1371/journal.pone.0199345>.
 46. Yilmaztepe MA, Yilmaz E. Effect of transient ulnar artery compression on radial artery diameter. *Exp Ther Med.* 2018;16(4):3735–9, <http://dx.doi.org/10.3892/etm.2018.6632>.
 47. Tsung JW, Blaivas M, Stone MB. Feasibility of point-of-care colour Doppler ultrasound confirmation of intraosseous needle placement during resuscitation. *Resuscitation.* 2009;80(6):665–8, <http://dx.doi.org/10.1016/j.resuscitation.2009.03.009>.
 48. Weekes AJ, Keller SM, Efuno B, Ghali S, Runyon M. Prospective comparison of ultrasound and CXR for confirmation of central vascular catheter placement. *Emerg Med J.* 2016;33(3):176–80, <http://dx.doi.org/10.1136/emermed-2015-205000>.
 49. Blanco P, Figueroa L, Menéndez MF, Berrueta B. Pericardiocentesis: ultrasound guidance is essential. *Ultrasound J.* 2022;14(1):9, <http://dx.doi.org/10.1186/s13089-022-00259-5>.
 50. Blanco P, Volpicelli G. Looking a bit superficial to the pleura. *Crit Ultrasound J.* 2014;6(1):13, <http://dx.doi.org/10.1186/s13089-014-0013-1>.
 51. Blanco P, Bello G. Agitated saline echo testing: two useful indications in critical care patients. *Med Intensiva.* 2019;43(2):124–5, <http://dx.doi.org/10.1016/j.medin.2018.09.016>.
 52. Maslove DM, Chen BT, Wang H, Kuschner WG. The diagnosis and management of pleural effusions in the ICU. *J Intensive Care Med.* 2013;28(1):24–36, <http://dx.doi.org/10.1177/0885066611403264>.
 53. Dancel R, Schnobrich D, Puri N, Franco-Sadud R, Cho J, Griksis L, et al. Recommendations on the use of ultrasound guidance for adult thoracentesis: a position statement of the Society of Hospital Medicine. *J Hosp Med.* 2018;13(2):126–35, <http://dx.doi.org/10.12788/jhm.2940>.
 54. Mercaldi CJ, Lanes SF. Ultrasound guidance decreases complications and improves the cost of care among patients undergoing thoracentesis and paracentesis. *Chest.* 2013;143(2):532–8, <http://dx.doi.org/10.1378/chest.12-0447>.
 55. Havelock T, Teoh R, Laws D, Gleeson F, BTS Pleural Disease Guideline Group. Pleural procedures and thoracic ultrasound: British Thoracic Society Pleural Disease Guideline 2010. *Thorax.* 2010;65 Suppl 2:i61–76, <http://dx.doi.org/10.1136/thx.2010.137026>.
 56. Millington SJ, Koenig S. Better with ultrasound: pleural procedures in critically ill patients. *Chest.* 2018;153(1):224–32, <http://dx.doi.org/10.1016/j.chest.2017.06.043>.
 57. Jayakrishnan B, Kashoob M, Al-Sukaiti R, Al-Mubaihsi S, Kakaria A, Al-Ghafri A, et al. Percutaneous ultrasound-guided pigtail catheter for pleural effusions: efficacy and safety. *Oman Med J.* 2021;36(2):e248, <http://dx.doi.org/10.5001/omj.2021.19>.
 58. Singh K, Loo S, Bellomo R. Pleural drainage using central venous catheters. *Crit Care.* 2003;7(6):R191–194, <http://dx.doi.org/10.1186/cc2393>.
 59. Volpicelli G, Boero E, Stefanone V, Storti E. Unusual new signs of pneumothorax at lung ultrasound. *Crit Ultrasound J.* 2013;5(1):10, <http://dx.doi.org/10.1186/2036-7902-5-10>.
 60. Blanco P, Figueroa L, Menéndez MF. Hydro-pneumothorax diagnosed by lung ultrasound in a COVID-19 patient. *Med Intensiva.* 2021;(Sep 17), <http://dx.doi.org/10.1016/j.medin.2021.08.009>. Epub ahead of print.
 61. Volpicelli G, Lamorte A, Tullio M, Boero E, Stefanone V. Worsening dyspnea and cough following thoracentesis. *Chest.* 2013;144(2):e1–3, <http://dx.doi.org/10.1378/chest.13-0674>.
 62. Cho J, Jensen TP, Reiersen K, Mathews BK, Bhagra A, Franco-Sadud R, et al. Recommendations on the use of ultrasound guidance for adult abdominal paracentesis: a position statement of the Society of Hospital Medicine. *J Hosp Med.* 2019;14:E7–15, <http://dx.doi.org/10.12788/jhm.3095>.
 63. Millington SJ, Koenig S. Better with ultrasound: paracentesis. *Chest.* 2018;154(1):177–84, <http://dx.doi.org/10.1016/j.chest.2018.03.034>.
 64. Agah S, Tavakoli S, Nikbakht H, Najafi M, Al-Agha A. Central venous pressure catheter for large-volume paracentesis in refractory ascites. *Indian J Gastroenterol.* 2014;33(4):310–5, <http://dx.doi.org/10.1007/s12664-014-0448-0>.
 65. Barbic D, Chenkin J, Cho DD, Jelic T, Scheuermeyer FX. In patients presenting to the emergency department with skin and soft tissue infections what is the diagnostic accuracy of point-of-care ultrasonography for the diagnosis of abscess compared to the current standard of care? A systematic review and meta-analysis. *BMJ Open.* 2017;7(1):e013688, <http://dx.doi.org/10.1136/bmjopen-2016-013688>. Erratum in: *BMJ Open.* 2017;7(9):e013688corr1.
 66. O'Rourke K, Kibbee N, Stubbs A. Ultrasound for the evaluation of skin and soft tissue infections. *Mo Med.* 2015;112(3):202–5. Retrieved December 22, 2022 from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6170135/>
 67. Gaspari RJ, Resop D, Mendoza M, Kang T, Blehar D. A randomized controlled trial of incision and drainage versus ultrasonographically guided needle aspiration for skin abscesses and the effect of methicillin-resistant *Staphylococcus aureus*. *Ann Emerg Med.* 2011;57(5):483–91, <http://dx.doi.org/10.1016/j.annemergmed.2010.11.021>, e1.
 68. Gaspari RJ, Sanseverino A, Gleeson T. Abscess incision and drainage with or without ultrasonography: a randomized controlled trial. *Ann Emerg Med.* 2019;73(1):1–7, <http://dx.doi.org/10.1016/j.annemergmed.2018.05.014>.
 69. Blanco P, Menéndez MF, Figueroa L, Provasi J, Blaivas M. Ultrasound-aided diagnosis of septic arthritis of the elbow in the emergency department. *J Ultrasound.* 2022;25(2):315–8, <http://dx.doi.org/10.1007/s40477-020-00506-2>.
 70. Wu T, Dong Y, Song HX, Fu Y, Li JH. Ultrasound-guided versus landmark in knee arthrocentesis: a systematic review. *Semin Arthritis Rheum.* 2016;45(5):627–32, <http://dx.doi.org/10.1016/j.semarthrit.2015.10.011>.
 71. Rowbotham EL, Grainger AJ. Ultrasound-guided intervention around the hip joint. *AJR Am J Roentgenol.* 2011;197(1):W122–127, <http://dx.doi.org/10.2214/ajr.10.6344>.

72. Berona K, Abdi A, Menchine M, Mailhot T, Kang T, Seif D, et al. Success of ultrasound-guided versus landmark-guided arthrocentesis of hip, ankle, and wrist in a cadaver model. *Am J Emerg Med.* 2017;35(2):240–4, <http://dx.doi.org/10.1016/j.ajem.2016.10.056>.
73. Gibbons RC, Zanaboni A, Genninger J, Costantino TG. Ultrasound-versus landmark-guided medium-sized joint arthrocentesis: a randomized clinical trial. *Acad Emerg Med.* 2022;29(2):159–63, <http://dx.doi.org/10.1111/acem.14396>.
74. Thom C, Ahmed A, Kongkatong M, Moak J. Point-of-care hip ultrasound leads to expedited results in emergency department patients with suspected septic arthritis. *J Am Coll Emerg Physicians Open.* 2020;1(4):512–20, <http://dx.doi.org/10.1002/emp2.12167>.
75. Soni NJ, Franco-Sadud R, Schnobrich D, Dancel R, Tierney DM, Salame G, et al. Ultrasound guidance for lumbar puncture. *Neurol Clin Pract.* 2016;6(4):358–68, <http://dx.doi.org/10.1212/cpj.0000000000000265>.
76. Shaikh F, Brzezinski J, Alexander S, Arzola C, Carvalho JC, Beyene J, et al. Ultrasound imaging for lumbar punctures and epidural catheterisations: systematic review and meta-analysis. *BMJ.* 2013;346:f1720, <http://dx.doi.org/10.1136/bmj.f1720>.
77. Soni NJ, Franco-Sadud R, Kobaidze K, Schnobrich D, Salame G, Lenchus J, et al. Recommendations on the use of ultrasound guidance for adult lumbar puncture: a position statement of the Society of Hospital Medicine. *J Hosp Med.* 2019;14(10):591–601, <http://dx.doi.org/10.12788/jhm.3197>.
78. Ravi PR, Naik S, Joshi MC, Singh S. Real-time ultrasound-guided spinal anaesthesia vs pre-procedural ultrasound-guided spinal anaesthesia in obese patients. *Indian J Anaesth.* 2021;65(5):356–61, http://dx.doi.org/10.4103/ija.ija_446_20.
79. Kusulas MP, Eutsler EP, DePiero AD. Bedside ultrasound for the evaluation of epidural hematoma after infant lumbar puncture. *Pediatr Emerg Care.* 2020;36(9):e508–12, <http://dx.doi.org/10.1097/pec.0000000000001383>.
80. Convissar D, Bittner EA, Chang MG. Biplane imaging using portable ultrasound devices for vascular access. *Cureus.* 2021;13(1):e12561, <http://dx.doi.org/10.7759/cureus.12561>.