



IMAGES IN INTENSIVE MEDICINE

Biphasic transdiaphragmatic pressure morphology identifies unilateral diaphragmatic paralysis



La presión transdiafrágica bifásica identifica parálisis diafrágica unilateral

Manuel Valdivia Marchal, José Ricardo Naranjo Izurieta, José Manuel Serrano Simón*

Servicio de Medicina Intensiva, Hospital Regional Universitario Reina Sofía, Córdoba, Spain

Available online 28 May 2024

A 60-year-old patient with a history of emphysema who failed to wean from mechanical ventilation seven days after right-unilateral lung transplantation, an elevation of the left hemi-diaphragm was noticed (Fig. 1). To evaluate respiratory muscle mechanics transdiaphragmatic pressure (Pdi) was monitored during pressure support, revealing insufficient respiratory muscle strength showed in Pdi and $\Delta P_{gas}/\Delta P_{es}$ ratio (Fig. 2, upper panel); as well in calculate partitioning pressure-time product (PTP) (Fig. 3). A remarkable biphasic morphology of Pdi during the inspiration was observed (Fig. 2, bottom panel), explained by the negative intrathoracic pressure drags the paretic and subsequent twitch of the competent hemi-diaphragm (arrows). This biphasic morphology of Pdi can distinguish hemi-diaphragm paralysis from other diaphragm dysfunction. The patient needs home ventilation.




Figure 1

* Corresponding author.

E-mail address: jm.serranosimon@gmail.com
(J.M. Serrano Simón).

<https://doi.org/10.1016/j.medine.2024.05.010>

2173-5727/© 2024 Elsevier España, S.L.U. and SEMICYUC. All rights are reserved, including those for text and data mining, AI training, and similar technologies.




Figure 2




Figure 3

Funding

No funding has been obtained for this work.

Conflict of interest

We declare that we have no conflict of interest.

Author contributions

MVM collected and analyzed the patient data and revised the manuscript. JRNI analyzed the patient data and revised the

manuscript. JMSS designed the study, interpreted all data and wrote the manuscript.

Acknowledgments

Dr. Robles Arista JC, and Dra Carmona Sánchez P contributed to the recruitment of patients. Dra De La Fuente Martos MC in encouragement to finish this manuscript.