



ORIGINAL

Evaluation of two intensive care models in relation to successful extubation after cardiac surgery



G.-W. Hao^{a,1}, G.-G. Ma^{a,1}, B.-F. Liu^{b,1}, X.-M. Yang^a, D.-M. Zhu^a, L. Liu^a, Y. Zhang^a, H. Liu^a, Y.-M. Zhuang^a, Z. Luo^a, G.-W. Tu^{a,*}

^a Department of Critical Care Medicine, Zhongshan Hospital, Fudan University, No. 180 Fenglin Road, Xuhui District, Shanghai 200032, PR China

^b Department of Critical Care Medicine, The First People's Hospital of Zhangjiagang, Suzhou, China

Received 19 April 2018; accepted 2 July 2018

Available online 23 August 2018

KEYWORDS

Extubation;
Cardiac surgery;
Care delivery;
Mechanical
ventilation

Abstract

Objective: To compare outcomes between intensivist-directed and cardiac surgeon-directed care delivery models.

Design: This retrospective, historical-control study was performed in a cohort of adult cardiac surgical patients at Zhongshan Hospital (Fudan University, China). During the first phase (March to August 2015), cardiac surgeons were in charge of postoperative care while intensivists were in charge during the second phase (September 2015–June 2016). Both phases were compared regarding successful extubation rate, intensive care unit (ICU) length of stay (LOS), and in-hospital mortality.

Setting: Tertiary Zhongshan Hospital (Fudan University, China).

Patients: Consecutive adult patients admitted to the cardiac surgical ICU (CSICU) after heart surgery.

Interventions: Phase I patients treated by cardiac surgeons, and phase II patients treated by intensivists.

Main variables of interest: Successful extubation, ICU LOS and in-hospital mortality.

Results: A total of 1792 (phase I) and 3007 patients (phase II) were enrolled. Most variables did not differ significantly between the two phases. However, patients in phase II had a higher successful extubation rate (99.17% vs. 98.55%; $p=0.043$) and a shorter median duration of mechanical ventilation (MV) (18 vs. 19 h; $p<0.001$). In relation to patients with MV duration >48 h, those in phase II had a comparatively higher successful extubation rate ($p=0.033$), shorter ICU LOS ($p=0.038$) and a significant decrease in in-hospital mortality ($p=0.039$).

* Corresponding author.

E-mail addresses: luo.zhe@zs-hospital.sh.cn (Z. Luo), tu.guowei@zs-hospital.sh.cn (G.-W. Tu).

¹ Guang-wei Hao, Guo-guang Ma and Bo-fei Liu contributed equally to this article.

PALABRAS CLAVE

Extubación;
Cirugía cardíaca;
Manejo médico;
Ventilación mecánica

Conclusions: The intensivist-directed care model showed improved rates of successful extubation and shorter MV durations after cardiac surgery.

© 2018 Published by Elsevier España, S.L.U.

Evaluación de 2 modelos de cuidados intensivos en cuanto a extubación exitosa de pacientes después de cirugía cardíaca

Resumen

Objetivo: Comparar el manejo entre intensivistas y cirujanos de pacientes de cirugía cardíaca en la unidad de cuidados intensivos.

Diseño: Este estudio de control retrospectivo se llevó a cabo con una cohorte de pacientes adultos de cirugía cardíaca. Durante la primera fase (de marzo a agosto de 2015), los cirujanos cardíacos estuvieron a cargo del manejo médico en la unidad; y durante la segunda fase (septiembre de 2015 a junio de 2016), lo hicieron intensivistas. Comparamos las fases en cuanto al número de extubaciones exitosas, el tiempo de estancia y la mortalidad.

Lugar del estudio: Hospital Zhongshan de la Universidad de Fudan, China.

Pacientes: Adultos admitidos secuencialmente a la unidad de cuidados intensivos cardíacos después de intervenciones quirúrgicas.

Intervenciones: Manejo médico por cirujanos en la primera fase, y por intensivistas en la segunda fase.

Variables prioritarias: Extubación exitosa, tiempo de estancia en la unidad de cuidados intensivos y mortalidad.

Resultados: Participaron 1.792 pacientes en la fase I y 3.007 en la fase II. Los pacientes de la fase II tuvieron más extubaciones exitosas (99,17 frente al 98,55%, $p=0,043$), y necesitaron menos tiempo de ventilación mecánica (mediana de 18 frente a 19 h, $p<0,001$). De entre los pacientes con ventilación mecánica de más de 48 h, los de la fase II fueron extubados exitosamente más veces, tuvieron una estancia más corta ($p=0,038$), y una menor mortalidad ($p=0,039$).

Conclusiones: El manejo médico por intensivistas aumentó significativamente el número de extubaciones exitosas y disminuyó el tiempo de ventilación mecánica.

© 2018 Publicado por Elsevier España, S.L.U.

Introduction

Cardiac surgery is now frequently offered to patients with advanced age, diabetes mellitus, chronic obstructive pulmonary disease, higher European system for cardiac operative risk evaluation II (EuroSCORE II) scores, and lower left ventricular ejection fractions, etc. Thus, the comorbidities in cardiac surgical patients have made the cases more complex over the past two decades.¹⁻⁴ This demographic change has resulted in a higher incidence of perioperative complications, prolonged intensive care unit (ICU) and hospital lengths of stay (LOS), and even hospital mortalities.⁵⁻⁹ As a result, the management of cardiac surgical patients requires a new level of critical care performance with introduction of dedicated and specialized critical care physicians.

Although the American Heart Association has recommended modifications to the staffing model for cardiac patients' care units,¹⁰ substantial variability remains in the organization for cardiac surgical intensive care units (CSICU).¹¹ In China, most CSICUs are affiliated to the department of cardiac surgery and are directed by cardiac surgeons. But the intensivists may have a role in many

aspects of patient care, particularly in improving ventilatory support, and the role of cardiac surgeons in CSICUs may be reduced in the future. Studies have evaluated the role of intensivists on patient outcomes after cardiac surgery, but have focused on mortality and ICU LOS.¹²⁻¹⁴ Few studies have paid attention to the ratio of successful extubations and no one has tried to demonstrate the reasons for extubation failure in adult cardiac surgical patients. The aim of this study was to assess whether an intensivist-directed care delivery model could improve the ratio of successful extubation after cardiac surgery and decrease ICU LOS and in-hospital mortality.

Patients and methods**Study design**

This single-center, retrospective, historical-control study was performed in a cohort of adult patients who underwent cardiac surgery at the Zhongshan Hospital of the Fudan University from March 2015 to June 2016. This is an academic teaching hospital with more than 3000

cardiac surgical procedures per year. In September 2015, the management of CSICU in our hospital was transferred from the department of cardiac surgery to the department of critical care medicine for quality improvement. Thus, a different care delivery model directed by intensivists was adopted. Accordingly, we collected clinical data of patients in the CSICU from March to August 2015 for a phase I group and from September 2015 to June 2016 for a phase II group. In phase I, cardiac surgeons who spent most of their time in the operating room directed the care delivery. Therapy decisions including extubation and discharges from the CSICU were often made before they got into the operation room. The cardiac surgeons managed the patients principally according to their personal experience, with consultant physicians from other departments including radiology, ultrasonography, and respiratory departments. In phase II, intensivists directed the care delivery and were 24 h/7 days on duty in the CSICU. Physicians from other departments supported the intensivists and specialized techniques (point of care [POC] ultrasound, fiberoptic scope, and advanced hemodynamic monitoring) were available to them. All intensivists extubated the patients according to the same criteria: clear consciousness, stable hemodynamics, adequate oxygenation, and successful spontaneous breathing trial. The spontaneous breathing trial was carried out using a continuous positive airway pressure or pressure support model, with pressure support at 5 cm H₂O and positive end expiratory pressures at 5 cm H₂O, lasting 30–60 min. The patients passed the spontaneous breathing trial if none of the following criteria were present: breathing frequency >35 breaths/min, SpO₂ <90%, rapid shallow breathing index (respiratory rate/tidal volume) >105 breaths/min/L, 20% increase or decrease from the baseline heart rate or blood pressure, use of accessory muscles, abdominal paradox movement, substantial agitation, anxiety, and/or diaphoresis.^{15,16} If the patient was not able to pass the spontaneous breathing trial, the intensivists tried to detect the reason for the failure using disciplinary related practices. The intensivists followed the directive protocol strictly regarding the timing of patient transferring to the ward: (1) weaning from various life support techniques; (2) respiratory and hemodynamic stabilities; (3) lack of severe discomfort feelings.

The primary outcome of this study was the improvement in successful extubations defined as not requiring reintubation within the first 48 h after the extubation. The secondary outcomes were the improvement of in-hospital mortality and the reduced ICU LOS.

Population

All consecutive patients older than 18 years, who underwent cardiac surgery between March 2015 and June 2016, were enrolled in this study. We excluded patients who did not undergo surgical treatment, those who died within 24 h after surgery, and those readmitted to the CSICU. The Ethical Committee of Zhongshan hospital affiliated to the Fudan University approved this study (NO. B2018-011). Informed consent forms were not required because this retrospective study did not modify existing diagnostic or therapeutic strategies.

Definitions

Extubation failure was asserted in cases needing reintubation within 48 h after extubation. Physicians identified atelectasis and pulmonary edema based on clinical manifestations, chest radiographies and/or ultrasound examinations. Stroke was defined as brain hemorrhage or infarction confirmed by radiographic examinations. CO₂ retention was identified in cases with high arterial blood CO₂ partial pressure and lethargy. The MV duration was defined as the time between ICU admission and the first extubation.

Data collection

The following data were recorded:

- (1) Baseline demographic data (age, gender, body mass index).
- (2) Patients' degree of disease severity (Acute Physiology and Chronic Health Evaluation II [APACHE II] score and EuroSCORE).
- (3) Type of surgery (valve, coronary artery bypass graft [CABG], valve and CABG, aortic, congenital and others).
- (4) Postoperative characteristics (renal replacement therapy [RRT], intra-aortic balloon pump [IABP], extracorporeal membrane oxygenation [ECMO], POC ultrasound, fiberoptic scope examination, computed tomography [CT] scan, and Pulse Contour Cardiac Output monitoring system [PiCCO]).
- (5) Outcome (MV duration, tracheotomy, noninvasive ventilation, successful extubation, ICU LOS, readmission, and in-hospital mortality).

Statistical methods

Continuous variables were presented as means ± SD or medians (25%–75% interquartile ranges, IQRs), while categorical variables were reported as adjusted proportions. Continuous data were compared using student *t*-test or the Mann–Whitney *U* test as appropriate, while differences between categorical variables were compared using the chi-square test or Fisher's exact test when necessary. A *p* value <0.05 was considered statistically significant. Statistical analyses were performed using the stata13.0 and SPSS 22.0 software (IBM Corporation, NY, USA).

Results

Demographic and clinical characteristics of enrolled patients

A total of 4799 patients were enrolled in this study. A patient flowchart was shown in Fig. 1. The patients enrolled in the phase I and phase II groups were 1792 and 3007, respectively. Demographic characteristics, APACHE II score, EuroSCORE, surgery type, postoperative characteristics, and outcomes were shown in Table 1. We found no significant differences between the two phases in terms of age, APACHE II score, EuroSCORE or type of surgery (5 subgroups: valve only, CABG only, valve and CABG, aortic, congenital




Figure 1 Flow diagram of the study.

and others). The cardiac surgeons did not perform POC ultrasound and fiberoptic scope examinations themselves, nor did they perform advanced hemodynamic monitoring. In phase I, the cardiac surgeons took 33 out of 1792 (1.8%) patients for CT scans; while in phase II, the intensivists took 118 out of 3007 (3.9%) patients for CT scans.

Outcome of enrolled patients

In phase II, patients had a higher ratio of successful extubation (99.2%) than in phase I (98.6%), $p=0.043$, together with a shorter median MV time (18 vs. 19 h, $p<0.001$). The differences in in-hospital mortality between the phase II (1.4%) and phase I (2.1%) and ICU LOS (39 vs. 38 h) were not significant. However, the readmission rate in phase II was lower (3.2%) than in phase I (4.7%), $p=0.008$. ECMO, RRT or IABP therapies were occasionally used in both phases. Tracheotomy (0.9% vs. 1.1%) and noninvasive ventilation (4.8% vs. 4.5%) were comparable in the two phases as well (Table 1).

Characteristics of the patients with extubation failure

In phase I, 26 out of 1792 patients were reintubated within 48 h after extubation, and in phase II, 25 out of 3007 patients were reintubated within 48 h after extubation. Demographic characteristics were comparable between the two phases. There were also no significant differences in terms of the type of surgery, tracheotomy and mortality between the two phases (Table 2). We found significant differences between the two phases concerning the causes of extubation failure (Table 3).

Characteristics of the patients with MV duration longer than 48 h

In phase I, 141 out of 1792 patients (7.9%) received MV for more than 48 h whilst in phase II, 245 out of 3007 patients (8.1%) underwent MV for longer than 48 h. We found no

significant differences in the two phases in terms of age, APACHE II, EuroSCORE and type of surgery. The two phases had patients undergoing similar life support techniques such as RRT, IABP and ECMO. In phase II, the intensivists performed more POC ultrasound and fiberoptic scope examinations, and they also took more patients for CT scans. We found no significant differences between the two phases in terms of MV durations, NIV or tracheotomy. Compared with phase I, phase II had a significantly higher ratio of successful extubations, decreasing in-hospital mortality, readmission rate, and lower ICU LOS (Table 4).

Discussion

In this historical cohort study comparing two different care delivery models in our dedicated CSICU, we found a significant improvement in the ratio of successful extubations in the intensivist-directed care delivery model. Additionally, the MV duration was shorter compared with that in the cardiac surgeon-directed model. The main reason for extubation failure in the phase I group was the presence of adverse respiratory variables such as secretions and atelectasis, while in phase II it was the presence of adverse cardiac variables. Other important clinical outcomes such as ICU LOS and in-hospital mortality were similar between the two groups. However, for the patients whose MV lasted >48 h, the ICU LOS and in-hospital mortality were both significantly better in the intensivist-directed care delivery model.

The rate of successful extubations is an important index to evaluate the quality of an ICU. This is especially true in the CSICU because almost all the cardiac surgical patients are admitted with mechanical ventilation postoperatively.¹⁷⁻¹⁹ To successfully extubate a patient, the physicians need to assess not only the pulmonary function but also other organ performances. The main training processes for intensivists and cardiac surgeons are quite different. The intensivists focus mainly on organ support techniques, while the cardiac surgeons care more about surgical related issues. Aging populations present more complicated situations and cardiothoracic critical care has become a new subspecialty requiring a specific curriculum

Table 1 Characteristics of patients in the two phase-groups.

Variables	Phase I (n = 1792)	Phase II (n = 3007)	p-Value
<i>Preoperative characteristics</i>			
Age (years)	59 ± 14	59 ± 16	0.547
Gender			0.947
Male (n)	1050	1759	
Female (n)	742	1248	
BMI	22.95 ± 3.23	23.44 ± 3.65	0.395
APACHE II score	7 ± 2	7 ± 3	0.754
EuroSCORE	4 ± 2	4 ± 2	0.763
Type of surgery, n (%)			0.655
Valve only	838 (46.8)	1394 (46.4)	
CABG only	302 (16.9)	539 (17.9)	
Valve and CABG	68 (3.8)	110 (3.7)	
Aortic	139 (7.8)	253 (8.4)	
Congenital	72 (4.0)	133 (4.4)	
Others	373 (20.8)	578 (19.2)	
<i>Postoperative characteristics</i>			
RRT, n (%)	59 (3.3)	91 (3.0)	0.608
IABP, n (%)	12 (0.7)	26 (0.9)	0.505
ECMO, n (%)	5 (0.3)	14 (0.5)	0.355
POC ultrasound, n (%)	0 (0)	2368 (78.8)	0
Fibreopticscopy, n (%)	0 (0)	216 (7.2)	0
CT scan, n (%)	33 (1.8)	118 (3.9)	0
PiCCO, n (%)	0 (0)	23 (0.8)	0
<i>Outcome</i>			
MV time, h	19 (8.24)	18 (17.20)	<0.001
Tracheotomy, n (%)	20 (1.1)	28 (0.9)	0.533
NIV, n (%)	80 (4.5)	143 (4.8)	0.643
Successful extubation, n (%)	1766 (98.6)	2982 (99.2)	0.043
ICU LOS, h	38 (22.70)	39 (23.72)	0.094
Readmission, n (%)	84 (4.7)	96 (3.2)	0.008
In hospital mortality, n (%)	37 (2.1)	43 (1.4)	0.097

Abbreviations: BMI, body mass index; APACHE, Acute Physiology and Chronic Health Evaluation; EuroSCORE, European system for cardiac operative risk evaluation; CABG, coronary artery bypass graft; RRT, renal replacement therapy; IABP, intra-aortic balloon pump; ECMO, extracorporeal membrane oxygenation; POC, point of care; CT, computed tomography; PiCCO, Pulse Contour Cardiac Output monitoring system; MV, mechanical ventilation; NIV, non-invasive ventilation; LOS, length of stay.

and ICU experience, which is not a specialty of surgical trainees.^{20–22} Intensivists perform a more thorough systematic evaluation of the patients and, thus, monitor important variables affecting successful extubation. Studies have shown that systematically evaluating risk factors other than pulmonary mechanics may reduce extubation failure.²³ To the best of our knowledge, this is the first study to assess the influence of different care delivery models on the ratio of successful extubation in adult cardiac surgical patients.

In this study, the intensivist-directed care delivery model not only improved the rate of successful extubations but also reduced the MV duration. This may be attributed to the different care delivery models. Although the reason for shorter mechanical ventilation periods during the phase II remained undetermined, our findings were consistent with those of previous studies.^{24–26} We hypothesize that perhaps it was the intensivists' presence at the bedside being able to diagnose and treat patients more quickly. And, we suggest that the intensivist-directed care delivery model may facilitate earlier extubations. Patients in phase II received more

advanced hemodynamic monitoring and POC examinations, such as bedside ultrasound examination. POC ultrasound can help to quickly identify the causes of weaning failure such as pulmonary edema,²⁷ pneumothorax, pulmonary atelectasis, and even diaphragmatic dysfunction.²⁸ And, advanced hemodynamical monitoring such as PiCCO also helps in guiding fluid resuscitation, and may lead to earlier extubations in hemodynamic compromised patients.²⁹

The differences of other major clinical outcomes such as in-hospital mortality and ICU LOS were not significant between phase I and phase II patients. This is consistent with studies regarding the influence of transition to a 24/7 in-house intensivist care delivery model on postoperative outcomes.^{13,26} However, other studies have shown that staffing ICUs with intensivists may improve clinical outcomes such as mortality and ICU LOS.^{30–32} The mortality and ICU LOS are influenced by many factors, including the population enrolled. More than 90% of our patients were extubated within 48 h after the operation. This group of patients displayed very low mortality and were often

Table 2 Characteristics of patients with extubation failure.

Variables	Phase I (n = 26)	Phase II (n = 25)	p-Value
Preoperative characteristics			
Age (years)	58 ± 16	62 ± 13	0.506
Gender			0.895
Male (n)	14	13	
Female (n)	12	12	
Smoking, n (%)	6 (23.1)	5 (20.00)	0.789
COPD, n (%)	3 (11.5)	2 (8.00)	0.671
Diabetes, n (%)	7 (26.9)	6 (24.00)	0.811
Hypertension, n (%)	13 (50.00)	12 (48.00)	0.886
History of cardiac surgery, n (%)	2 (7.7)	4 (16.00)	0.357
Preoperative Hb, g/L	131.83 ± 15.19	123.13 ± 24.49	0.17
Preoperative SCr, μmol/L	84.06 ± 24.16	104.53 ± 37.53	0.624
Preoperative EF, %	57 ± 10	54 ± 10	0.832
NYHA, III–IV, n (%)	10 (38.5)	9 (36.00)	0.856
APACHE II score	14 ± 5	15 ± 7	0.028
EuroSCORE	6 ± 3	6 ± 4	0.384
Operative characteristics			
Type of surgery			0.915
Valve only, n (%)	9 (34.6)	8 (32.00)	
CABG only, n (%)	6 (23.1)	5 (20.00)	
Valve + CABG, n (%)	4 (15.4)	7 (28.00)	
Aortic, n (%)	3 (11.5)	2 (8.00)	
Congenital, n (%)	2 (7.7)	1 (4.00)	
Others, n (%)	2 (7.7)	2 (8.00)	
Emergency, n (%)	0 (0)	1 (4.00)	0.303
Outcome			
Tracheotomy, n (%)	9 (34.6)	7 (28.00)	0.611
In hospital mortality, n (%)	4 (15.4)	3 (12.00)	0.725

Abbreviations: COPD, chronic obstructive pulmonary disease; Hb, hemoglobin; SCr, serum creatinine; EF, ejection fraction; NYHA, New York Heart Association; APACHE, Acute Physiology and Chronic Health Evaluation; EuroSCORE, European system for cardiac operative risk evaluation; CABG, coronary artery bypass graft.

Table 3 Reasons of extubation failure.

Variables	Phase I (n = 26)	Phase II (n = 25)	p-Value
Respiratory, n (%)			
Secretions	8	4	0.015
Bronchospasm	1	2	
Atelectasis	7	0	
Cardiac, n (%)	7 (26.9)	14 (56)	
Pulmonary edema	3	2	
Cardiac arrest	3	3	
Life threatening arrhythmias	1	9	
Neurological, n (%)			
Stroke and seizure	4 (15.4)	5 (20)	0.666
CO ₂ retention	2	3	
	2	2	

discharged within 24h after extubation, so their short ICU stay may not have provided enough time to show differences or benefits associated with the type of care received. Additionally, the shorter median MV duration in phase I among all patients was not clinically meaningful; and therefore, we reanalyzed our data in patients whose MV lasted >48h after the operation, and found that

reintubations, mortality and ICU LOS were significantly reduced in those in the phase II. Additionally, our data showed that patients with prolonged mechanical ventilation had higher APACHE II scores, and higher frequencies of IABP, RRT, PiCCO or ECMO usage. The predictors of prolonged mechanical ventilation after cardiac surgery include older age, cardiac dysfunction, chronic renal failure, chronic

Table 4 Characteristics of the patients with MV duration over 48 h.

Variables	Phase I (n = 141)	Phase II (n = 245)	p-Value
Preoperative characteristics			
Age (years)	58 ± 13	60 ± 14	0.126
Gender			0.239
Male (n)	97	154	
Female (n)	44	91	
BMI	22.96 ± 6.07	23.18 ± 5.67	0.82
APACHE II score	12 ± 6	13 ± 7	0.536
EuroSCORE	6 ± 2	6 ± 2	0.769
Type of surgery, n (%)			0.416
Valve only	53 (37.6)	119 (46.7)	
CABG only	26 (18.4)	50 (19.6)	
Valve + CABG	12 (8.5)	15 (5.9)	
Aortic only	7 (5.0)	13 (5.1)	
Congenital only	11 (7.8)	13 (5.1)	
Others	32 (22.7)	45 (17.7)	
Postoperative characteristics			
RRT, n (%)	39 (27.7)	55 (22.5)	0.251
IABP, n (%)	4 (2.8)	11 (4.5)	0.418
ECMO, n (%)	3 (2.1)	12 (4.9)	0.175
POC ultrasound, n (%)	0 (0)	209 (85.3)	0
Fibreopticscopy, n (%)	0 (0)	135 (55.1)	0
CT scan, n (%)	14 (9.9)	43 (17.6)	0.042
PiCCO, n (%)	0 (0)	22 (9.0)	0
Outcome			
MV time, h	91 (67,127)	84 (66,116)	0.126
Tracheotomy, n (%)	14 (9.93)	13 (5.31)	0.086
NIV, n (%)	27 (19.15)	62 (25.31)	0.167
Successful extubation, n (%)	126 (89.36)	233 (95.10)	0.033
ICU LOS, h	165 (97,391)	145 (100,221)	0.038
Readmission, n (%)	31 (21.99)	21 (8.57)	0
In hospital mortality, n (%)	21 (14.89)	20 (10.62)	0.039

Abbreviations: BMI, body mass index; APACHE, Acute Physiology and Chronic Health Evaluation; EuroSCORE, European system for cardiac operative risk evaluation; CABG, coronary artery bypass graft; RRT, renal replacement therapy; IABP, intra-aortic balloon pump; ECMO, extracorporeal membrane oxygenation; POC, point of care; CT, computed tomography; PiCCO, Pulse Contour Cardiac Output monitoring system; MV, mechanical ventilation; NIV, non-invasive ventilation; LOS, length of stay.

obstructive pulmonary disease, repeated surgery, emergency surgery, higher New York Heart Association/Canadian Cardiovascular Society class, longer cardiopulmonary bypass time, blood product transfusions,³³ early postoperative hemodynamic status, and events such as stroke and bacteremia.³⁴ In this study, the possible causes for prolonged mechanical ventilation were the presence of underlying chronic diseases before surgery, hemodynamic instability and surgery related complications. Management of these patients often required a broad view of both cardiac and other organ functions. The more comprehensive management offered by the intensivists may have resulted in the improved outcomes of the patients with prolonged mechanical ventilation in the second phase. Collectively, our results indicate that patients with prolonged mechanical ventilation may get more benefits from the intensivist-directed care delivery model.

The extubation failure may cause increased morbidity and longer ICU LOS.^{35,36} The main causes of extubation failure include respiratory, cardiac and neurological variables,

with respiratory failure being the most common cause.³⁷⁻³⁹ However, Forouzan et al. demonstrated that cardiac variables prompt more reintubations after cardiac surgery in adult patients than the respiratory variables.⁴⁰ We found significant differences on the causes of extubation failure between the two phase-groups. In phase I, 15 out of the 26 extubation failure cases (57.7%) were reintubated for respiratory reasons. But, respiratory variables such as secretions, bronchospasm, and atelectasis are reversible if managed properly. In phase II, only 6 out of the 25 (24%) extubation failure cases were reintubated for respiratory reasons. The majority of the reintubations were due to life-threatening arrhythmias and cardiac arrest, which typically cannot be prevented and are less amenable to treatment. We propose that intensivists are more skillful in airway management. Intensivists addressed most of the respiratory causes for extubation failure, and thus cardiac variables (56%) became the main cause of extubation failure. Our results indirectly indicate that the intensivist-directed care delivery model in our CSICU was superior to the cardiac

surgeon-directed model in terms of patient extubation management.

We are aware of the limitations of our study. First, this was a retrospective study, so we could not collect and analyze all clinical data details, such as results of POC ultrasound, hemodynamic monitoring and fiberoptic scope examinations. Second, whether the difference in care delivery models and the improved outcomes (reduced MV duration and higher ratio of successful extubation) had a cause-effect relationship remains to be confirmed. Finally, this was a single center study, and the results may not be applicable to other ICUs. Further prospective, multicenter trials are necessary to confirm the conclusions of this study.

Conclusions

Compared with the cardiac surgeon-directed care delivery model, the intensivist-directed model led to an improved successful extubation rate as well as a shorter duration of MV after cardiac surgery. The intensivist-directed care delivery model is superior and feasible in our CSICU, and should be considered for other CSICUs. Further research will be required before more definitive recommendations can be made.

Author's contributions

Guang-Wei Hao, Guo-Guang Ma and Bo-Fei Liu performed the literature search, extracted the data and drafted the manuscript. Xiao-Mei Yang, Ying Zhang, Lan Liu, Hua Liu and Ya-Min Zhuang reviewed studies for inclusion and extracted data. Guo-Wei Tu, Du-Ming Zhu and Zhe Luo performed the analysis and helped draft the manuscript. Guo-Wei Tu and Zhe Luo conceived the idea, participated in manuscript writing and revision. All authors have read and approved the final manuscript.

Funding

This article was supported by grants from the Natural Science Foundation of Shanghai (16ZR1405600), the National Natural Science Foundation of China (81500067), the Health and Family Planning Commission of Shanghai (20154Y011) and research funds of from the Zhong Shan Hospital (2017ZSYXQN23 and 2017ZSQN16).

Conflicts of interest

The authors declare that they have no conflicts of interest.

References

1. Ferguson TB Jr, Hammill BG, Peterson ED, DeLong ER, Grover FL. A decade of change – risk profiles and outcomes for isolated coronary artery bypass grafting procedures, 1990–1999: a report from the STS National Database Committee and the Duke Clinical Research Institute Society of Thoracic Surgeons. *Ann Thorac Surg.* 2002;73:480–9, discussion 89–90.
2. ElBardissi AW, Aranki SF, Sheng S, O'Brien SM, Greenberg CC, Gammie JS. Trends in isolated coronary artery bypass grafting: an analysis of the Society of Thoracic Surgeons adult cardiac surgery database. *J Thorac Cardiovasc Surg.* 2012;143:273–81.
3. Kindo M, Hoang Minh T, Perrier S, Bentz J, Mommerot A, Billaud P, et al. Trends in isolated coronary artery bypass grafting over the last decade. *Interact Cardiovasc Thorac Surg.* 2017;24:71–6.
4. Katz NM. The emerging specialty of cardiothoracic surgical critical care: the leadership role of cardiothoracic surgeons on the multidisciplinary team. *J Thorac Cardiovasc Surg.* 2007;134:1109–11.
5. Hein OV, Birnbaum J, Wernecke K, England M, Konertz W, Spies C. Prolonged intensive care unit stay in cardiac surgery: risk factors and long-term-survival. *Ann Thorac Surg.* 2006;81:880–5.
6. Atoui R, Ma F, Langlois Y, Morin JF. Risk factors for prolonged stay in the intensive care unit and on the ward after cardiac surgery. *J Card Surg.* 2008;23:99–106.
7. Shahian DM, O'Brien SM, Filardo G, Ferraris VA, Haan CK, Rich JB, et al. The Society of Thoracic Surgeons 2008 cardiac surgery risk models: part 1 – coronary artery bypass grafting surgery. *Ann Thorac Surg.* 2009;88 1 Suppl.:S2–22.
8. O'Brien SM, Shahian DM, Filardo G, Ferraris VA, Haan CK, Rich JB, et al. The Society of Thoracic Surgeons 2008 cardiac surgery risk models: part 2 – isolated valve surgery. *Ann Thorac Surg.* 2009;88 1 Suppl.:S23–42.
9. Rahmanian PB, Kroner A, Langebartels G, Ozel O, Wippermann J, Wahlers T. Impact of major non-cardiac complications on outcome following cardiac surgery procedures: logistic regression analysis in a very recent patient cohort. *Interact Cardiovasc Thorac Surg.* 2013;17:319–26, discussion 26–7.
10. Morrow DA, Fang JC, Fintel DJ, Granger CB, Katz JN, Kushner FG, et al. Evolution of critical care cardiology: transformation of the cardiovascular intensive care unit and the emerging need for new medical staffing and training models: a scientific statement from the American Heart Association. *Circulation.* 2012;126:1408–28.
11. Lane-Fall MB, Ramaswamy TS, Brown SES, He X, Gutsche JT, Fleisher LA, et al. Structural nursing, and physician characteristics and 30-day mortality for patients undergoing cardiac surgery in Pennsylvania. *Crit Care Med.* 2017;45:1472–80.
12. Kogan A, Preisman S, Berkenstadt H, Segal E, Kassif Y, Sternik L, et al. Evaluation of the impact of a quality improvement program and intensivist-directed ICU team on mortality after cardiac surgery. *J Cardiothorac Vasc Anesth.* 2013;27:1194–200.
13. Kumar K, Singal R, Manji RA, Zarychanski R, Bell DD, Freed DH, et al. The benefits of 24/7 in-house intensivist coverage for prolonged-stay cardiac surgery patients. *J Thorac Cardiovasc Surg.* 2014;148:290–7.e6.
14. Na SJ, Chung CR, Jeon K, Park CM, Suh GY, Ahn JH, et al. Association between presence of a cardiac intensivist and mortality in an adult cardiac care unit. *J Am Coll Cardiol.* 2016;68:2637–48.
15. Barbosa e Silva MG, Borges DL, Costa Mde A, Baldez TE, Silva LN, Oliveira RL, et al. Application of mechanical ventilation weaning predictors after elective cardiac surgery. *Braz J Cardiovasc Surg.* 2015;30:605–9.
16. Gupta P, Giehler K, Walters RW, Meyerink K, Modrykamien AM. The effect of a mechanical ventilation discontinuation protocol in patients with simple and difficult weaning: impact on clinical outcomes. *Respir Care.* 2014;59:170–7.
17. Legare JF, Hirsch GM, Buth KJ, MacDougall C, Sullivan JA. Preoperative prediction of prolonged mechanical ventilation following coronary artery bypass grafting. *Eur J Cardiothorac Surg.* 2001;20:930–6.
18. Meade MO, Guyatt G, Butler R, Elms B, Hand L, Ingram A, et al. Trials comparing early vs late extubation following cardiovascular surgery. *Chest.* 2001;120 6 Suppl.:445S–53S.
19. Filsoufi F, Rahmanian PB, Castillo JG, Chikwe J, Adams DH. Predictors and early and late outcomes of respiratory failure in contemporary cardiac surgery. *Chest.* 2008;133:713–21.

20. Katz NM. The evolution of cardiothoracic critical care. *J Thorac Cardiovasc Surg.* 2011;141:3–6.
21. Katz NM. It is time for certification in cardiothoracic critical care. *J Thorac Cardiovasc Surg.* 2013;145:1446–7.
22. Sherif HM. Developing a curriculum for cardiothoracic surgical critical care: impetus and goals. *J Thorac Cardiovasc Surg.* 2012;143:804–8.
23. Krinsley JS, Reddy PK, Iqbal A. What is the optimal rate of failed extubation? *Crit Care.* 2012;16:111.
24. Kumar K, Zarychanski R, Bell DD, Manji R, Zivot J, Menkis AH, et al. Impact of 24-hour in-house intensivists on a dedicated cardiac surgery intensive care unit. *Ann Thorac Surg.* 2009;88:1153–61.
25. Netzer G, Liu X, Shanholtz C, Harris A, Verceles A, Iwashyna TJ. Decreased mortality resulting from a multicomponent intervention in a tertiary care medical intensive care unit. *Crit Care Med.* 2011;39:284–93.
26. Benoit MA, Bagshaw SM, Norris CM, Zibdawi M, Chin WD, Ross DB, et al. Postoperative complications and outcomes associated with a transition to 24/7 intensivist management of cardiac surgery patients. *Crit Care Med.* 2017;45:993–1000.
27. Lamia B, Maizel J, Ochagavia A, Chemla D, Osman D, Richard C, et al. Echocardiographic diagnosis of pulmonary artery occlusion pressure elevation during weaning from mechanical ventilation. *Crit Care Med.* 2009;37:1696–701.
28. Dube BP, Dres M. Diaphragm dysfunction: diagnostic approaches and management strategies. *J Clin Med.* 2016;5.
29. Hendy A, Bubenek S. Pulse waveform hemodynamic monitoring devices: recent advances and the place in goal-directed therapy in cardiac surgical patients. *Rom J Anaesth Intensive Care.* 2016;23:55–65.
30. Vincent JL. Need for intensivists in intensive-care units. *Lancet.* 2000;356:695–6.
31. Pronovost PJ, Angus DC, Dorman T, Robinson KA, Dremsizov TT, Young TL. Physician staffing patterns and clinical outcomes in critically ill patients: a systematic review. *JAMA.* 2002;288:2151–62.
32. Sultan OW, Boland LL, Kinzy TG, Melamed RR, Seatter SC, Fari-var RS, et al. Improved outcomes with integrated intensivist consultation for cardiac surgery patients. *Am J Med Qual.* 2018, 1062860618766614.
33. Cislighi F, Condemni AM, Corona A. Predictors of prolonged mechanical ventilation in a cohort of 5123 cardiac surgical patients. *Eur J Anaesthesiol.* 2009;26:396–403.
34. Murthy SC, Arroliga AC, Walts PA, Feng J, Yared JP, Lytle BW, et al. Ventilatory dependency after cardiovascular surgery. *J Thorac Cardiovasc Surg.* 2007;134:484–90.
35. Gowardman JR, Huntington D, Whiting J. The effect of extubation failure on outcome in a multidisciplinary Australian intensive care unit. *Crit Care Resusc.* 2006;8:328–33.
36. Rady MY, Ryan T. Perioperative predictors of extubation failure and the effect on clinical outcome after cardiac surgery. *Crit Care Med.* 1999;27:340–7.
37. Kulkarni AP, Agarwal V. Extubation failure in intensive care unit: predictors and management. *Indian J Crit Care Med.* 2008;12:1–9.
38. Burns KE, Adhikari NK, Meade MO. A meta-analysis of non-invasive weaning to facilitate liberation from mechanical ventilation. *Can J Anaesth.* 2006;53:305–15.
39. Engoren M, Buderer NF, Zacharias A, Habib RH. Variables predicting reintubation after cardiac surgical procedures. *Ann Thorac Surg.* 1999;67:661–5.
40. Yazdanian F, Azarfari R, Aghdaii N, Faritous SZ, Motlagh SD, Panahpour A. Cardiac variables as main predictors of endotracheal reintubation rate after cardiac surgery. *J Tehran Heart Center.* 2013;8:42–7.