

- process of care and ICU structure to improved outcomes: an update from the American College of Critical Care Medicine Task Force on models of critical care. *Crit Care Med.* 2015;43:1520–5.
2. Chen LM, Render M, Sales A, Kennedy EH, Wiitala W, Hofer TP. Intensive care unit admitting patterns in the Veterans affairs health care system. *Arch Intern Med.* 2012;172:1220–6.
 3. Singer JP, Kohlwes J, Bent S, Zimmerman L, Eisner MD. The impact of a «low-intensity» versus «high-intensity» medical intensive care unit on patient outcomes in critically ill veterans. *J Intensive Care Med.* 2010;25:233–9.
 4. Lucena JF, Alegre F, Martinez-Urbistondo D, Landecho MF, Huerta A, García-Mouriz A, et al. Performance of SAPS II and SAPS 3 in intermediate care. *PLoS ONE.* 2013;8:e77229.
 5. Moreno RP, Metnitz PG, Almeida E, Jordan B, Bauer P, Campos RA, et al. SAPS 3 – From evaluation of the patient to evaluation of the intensive care unit. Part 2: Development of a prognostic model for hospital mortality at ICU admission. *Intensive Care Med.* 2005;31:1345–55.
 6. Chang DW, Shapiro MF. Association between intensive care unit utilization during hospitalization and costs, use of invasive procedures, and mortality. *JAMA Intern Med.* 2016;176:1492–9.
 7. Katsounas A, Kamacharova I, Tyczynski B, Eggebrecht H, Erbel R, Canbay A, et al. The predictive performance of the SAPS II and SAPS 3 scoring systems: a retrospective analysis. *J Crit Care.* 2016;33:180–5.
 8. Confalonieri M, Trevisan R, Demisar M, Lattuada L, Longo C, Cifaldi R, et al. Opening of a respiratory intermediate care unit in a general hospital: impact on mortality and other outcomes. *Respiration.* 2015;90:235–42.
 9. Edbrooke DL, Minelli C, Mills GH, Iapichino G, Pezzi A, Corbella D, et al. Implications of ICU triage decisions on patient mortality: a cost-effectiveness analysis. *Crit Care.* 2011;15:R56.
 10. Kahn JM, Rubenfeld GD, Rohrbach J, Fuchs BD. Cost savings attributable to reductions in intensive care unit length of stay for mechanically ventilated patients. *Med Care.* 2008;46:1226–33.
- R. Fernandez^{a,b,c,*}, J.M. Alcoverro^a, I. Catalán^a, O. Rubio^a, S. Cano^a, C. Subirà^a
- ^a *Servicio de Medicina Intensiva, Hospital Sant Joan de Déu-Fundació Althaia, Manresa, Barcelona, Spain*
^b *Universitat Internacional de Catalunya, Barcelona, Spain*
^c *CIBERES, Madrid, Spain*
- * Corresponding author.
 E-mail address: rfernandezf@althaia.cat (R. Fernandez).
 2173-5727/
 © 2017 Elsevier España, S.L.U. and SEMICYUC. All rights reserved.

Kidney transplantation in the Intensive Care Unit: Graft evaluation using imaging tests[☆]



Postoperatorio de trasplante renal en la unidad de cuidados intensivos: evaluación del injerto mediante técnicas de imagen

Dear Sir,

Kidney transplantation is the management of choice in patients with end-stage chronic kidney disease. For decades it has been shown to offer improved quality of life and a lesser mortality risk compared with hemodialysis.¹ However, transplantation remains a complex process and involves patients with important comorbidities. It is therefore essential to optimize the postoperative care of these individuals in the Intensive Care Unit (ICU). Imaging techniques play a very important role in this respect. Both renal doppler ultrasound (RDU) and radioisotopic studies (renal scintigraphy) are of great help in the identification of early complications, and afford very useful basal information for adequate graft assessment.^{2,3}

Renal doppler ultrasound performed at the patient bedside by the intensivist makes it possible to discard acute complications such as bleeding, hydronephrosis,

vascular dehiscence, arterial stenosis, venous thrombosis or renal infarction.² The technique is moreover inexpensive, noninvasive and avoids the need for patient transfer. Furthermore, the renal resistance index (RI) (RI = peak systolic flow – end-diastolic flow/peak systolic flow) allows the quantification of graft flow alterations. It was originally used to diagnose acute rejection,⁴ though recently it has also been regarded as a chronic rejection marker.⁵ On the other hand, RI > 0.8 has been shown to be a strong predictor of graft loss, being related to donor death.⁵ However, despite its prognostic value in transplant patients, RI is more closely related to recipient age and certain hemodynamic factors than to renal anomalies.^{6,7} Radioisotopic studies in turn allow evolutive monitoring of the graft. Although a single or point evaluation is unable to distinguish between acute tubular necrosis and acute rejection, serial radioisotopic studies evidencing a progressive decrease in function and perfusion could be indicative of acute rejection.⁸

In our ICU, renal doppler ultrasound is performed by the intensivist in the immediate postoperative period, while renal scintigraphy is routinely performed on the morning after the operation. In the event of an interval of more than 12 h between both explorations, a new ultrasound study is made before patient transfer. Since our Unit provides support for a large number of kidney transplants, we decided to conduct a retrospective observational study, analyzing the explorations made between January 2013 and December 2015 in all patients admitted to the ICU in the immediate postoperative period of kidney transplantation. In this context, on performing RDU, adequate flow was defined as RI ≤ 0.7 in the principal renal artery and in the segmental renal arteries. In relation to scintigraphy, mercaptoacetyl-glycine (MAG-3) is the radionuclide of choice in our center. The three phases of renal function were evaluated. The first phase shows graft perfusion based on images obtained during

[☆] Please cite this article as: Roldán-Reina AJ, Egea-Guerrero JJ, Palomo-López N, Cuenca-Apolo DX, Adriaenssens-Pérez M, Martín-Villén L. Postoperatorio de trasplante renal en la unidad de cuidados intensivos: evaluación del injerto mediante técnicas de imagen. *Med Intensiva.* 2019;43:384–386.

Table 1 General characteristics of the kidney transplant patients.

| Type of donor | Brain-dead donor | Type II non-heart beating donor | Type III non-heart beating donor | Live donor |
|---|------------------|---------------------------------|----------------------------------|------------------|
| <i>n</i> (%) | 189 (69.2) | 26 (9.5) | 17 (6.2) | 41 (15) |
| Lactate upon admission to ICU, median (IQR) [mmol/l] | 1.4 (1.0–2.0) | 1.4 (1.2–1.9) | 1.3 (1.1–1.7) | 1.5 (1.0–2.1) |
| Creatinine upon admission to ICU, median (IQR) [mg/dl] | 5.86 (4.45–7.23) | 6.83 (5.24–7.68) | 5.64 (4.78–6.62) | 5.5 (4.28–6.23) |
| Creatinine at hospital discharge, median (IQR) [mg/dl] ^a | 1.81 (1.36–2.48) | 4.26 (2.23–6.17) | 2.5 (2.21–3.96) | 1.47 (1.05–1.66) |
| Diuresis during the first 24 h in the ICU (%) | | | | |
| Polyuria | 58.0 | 38.5 | 41.2 | 92.7 |
| Oliguria | 31.4 | 42.3 | 35.3 | 7.3 |
| Anuria | 10.6 | 19.2 | 23.5 | 0.0 |

^a Statistically significant differences. IQR: interquartile range.

the first minute after intravenous injection of the tracer. The second phase evidences tracer extraction by the nephron and its excretion via glomerular filtration and/or tubular secretion. The third phase in turn assessed elimination of the tracer. Slight tubular dysfunction and/or a small delay in tracer elimination were considered normal.

A descriptive analysis was made, followed by comparison of the evolution of renal function in the transplant patients according to the type of donor involved. The Student's *t*-test was used in the presence of normal data distribution, while the Mann–Whitney *U*-test was applied in the absence of a normal distribution. The kappa test was used to assess agreement between the two flow tests used (RDU and scintigraphy). Statistical significance was considered for $p < 0.05$. The SPSS version 22.0 statistical package (IBM Corporation, NY, USA) was used throughout.

A total of 273 patients (60.1% males) were included during the study period. The mean age was 52 years. The patient characteristics are shown in Table 1, while agreement between the two tests is reported in Table 2 – the kappa index being 0.5 ($p < 0.001$).

Few studies to date have compared RDU versus radioisotopic explorations for the evaluation of early graft function. Cofán et al.⁹ found that RDU did not discriminate the severity of graft acute tubular necrosis, while altered renogram findings were correlated to an increased need for hemodialysis and a decrease in graft survival. Another study in pediatric patients¹⁰ found renal blood flow as calculated from radioisotopic techniques to offer a more reliable

assessment of graft dysfunction than RI measurement. A more recent study⁸ concluded that serial isotopic tests combined with kidney biopsy are more reliable in demonstrating graft dysfunction.

The great majority of specialized centers use both scintigraphy and RDU for evaluating the graft. However, in our study agreement (kappa index) with the perfusion indices was found to be moderate. Since most of the patients were admitted during hours on duty, RDU was performed by the intensivist available at the time. Although our team has extensive experience with transplants, the possible inter-observer effect of ultrasound could constitute a limitation in our study.

Kidney transplantation requires adequate postoperative care in order to guarantee good graft evolution in the first hours. In coincidence with other series, our results illustrate the early renal graft functional differences according to the type of donor involved. These differences are mainly due to the existence of longer warm ischemia times in the case of grafts harvested from type II non-heart beating donors. Nevertheless, it is known that these differences are minimized over the long term.¹¹ In this regard, RDU and scintigraphy contribute very valuable information and have made it possible to maintain good functional outcomes in recent years. Our findings suggest that it is advisable to continue using both diagnostic tools for evaluating kidney transplants, integrating the results of these two tests in the clinical context of each individual patient.

Conflicts of interest

The authors declare that they have no conflicts of interest.

References

1. Tennankore KK, Kim SJ, Baer HJ, Chan CT. Survival and hospitalization for intensive home hemodialysis compared with kidney transplantation. *J Am Soc Nephrol.* 2014;25:2113–20.
2. Nankivell BJ, Kuypers DR. Diagnosis and prevention of chronic kidney allograft loss. *Lancet.* 2011;378:1428–37.
3. Dubovsky EV, Russell CD, Vischof-Delaloye A, Bubeck B, Chaiwatanarat T, Hilson AJ, et al. Report of the radionuclides

Table 2 Agreement between the two tests (kappa index).

| Normal test | MAG-3, <i>n</i> (%) | | Total |
|-------------------|---------------------|----------|-------|
| | Yes | No | |
| <i>RDU, n</i> (%) | | | |
| Yes | 244 (94.6) | 14 (5.4) | 258 |
| No | 2 (18.2) | 9 (81.2) | 11 |
| Total | 246 | 23 | 269 |

RDU: renal doppler ultrasound; MAG-3: renal scintigraphy. Kappa index = 0.5 ($p < 0.001$).

- in Nephrourology Committee for Evaluation of Transplanted Kidney (review of techniques). *Semin Nucl Med.* 1999;29:175–88.
4. Rifkin MD, Needleman L, Pasto ME, Kurtz AB, Foy PM, McGlynn E, et al. Evaluation of renal transplant rejection by duplex Doppler examination: value of resistive index. *AJR Am J Roentgenol.* 1987;148:759–62.
 5. De Vries AP, van Son WJ, van der Heide JJ, Ploeg RJ, Navis G, de Jong PE, et al. The predictive value of renal vascular resistance for late renal allograft loss. *Am J Transplant.* 2006;6:364–70.
 6. Radermacher J, Mengel M, Ellis S, Stuht S, Hiss M, Schwarz A, et al. The renal arterial resistance index and renal allograft survival. *N Engl J Med.* 2003;349:115–24.
 7. Naesens M, Heylen L, Lerut E, Claes K, de Wever L, Claus F, et al. Intrarenal resistive index after renal transplantation. *N Engl J Med.* 2013;369:1797–806.
 8. Kocabaş B, Aktaş A, Aras M, Işıklar I, Gençoğlu A. Renal scintigraphy findings in allograft recipients with increased resistance index on Doppler sonography. *Transplant Proc.* 2008;40:100–3.
 9. Cofán F, Gilabert R, Oppenheimer F, Bru C, Lomeña F, Setoain F, et al. Duplex-Doppler ultrasound and MAG-3 scintigraphy in the evaluation of acute tubular necrosis after kidney transplantation. *Transplant Proc.* 1997;29:1376–7.
 10. Fitzpatrick MM, Gleeson FV, de Bruyn R, Trompeter RS, Gordon I. The evaluation of paediatric renal transplants using resistive index and renal blood flow. *Pediatr Nephrol.* 1992;6:172–5.
 11. Gentil MA, Castro de la Nuez P, Gonzalez-Corvillo C, de Gracia MC, Cabello M, Mazuecos MA, et al. Non-heart-beating donor kidney transplantation survival is similar to donation after brain death: comparative study with controls in a regional program. *Transplant Proc.* 2016;48:2867–70.
- A.J. Roldán-Reina^{a,*}, J.J. Egea-Guerrero^{a,b,c},
N. Palomo-López^a, D.X. Cuenca-Apolo^a,
M. Adriaenssens-Pérez^a, L. Martín-Villén^{a,b}
- ^a *UGC de Medicina Intensiva, Hospital Universitario Virgen del Rocío, Sevilla, Spain*
^b *Coordinación de Trasplantes, Hospital Universitario Virgen del Rocío, Sevilla, Spain*
^c *IBiS-CSIC-Universidad de Sevilla, Sevilla, Spain*
- * Corresponding author.
E-mail address: roldanyreina@gmail.com
(A.J. Roldán-Reina).
2173-5727/
© 2017 Elsevier España, S.L.U. and SEMICYUC. All rights reserved.