



Successive prone positioning sessions in mechanically ventilated patients with moderate and severe acute respiratory distress syndrome secondary to COVID-19: case series

Sucesivos ciclos de decúbito prono en pacientes bajo asistencia ventilatoria mecánica invasiva con distrés moderado a severo debido a COVID-19. Estudio descriptivo, tipo serie de casos

Dear Editor,

Prone positioning (PP) in patients with severe oxygenation impairment is one of the adjuvant strategies that has demonstrated to improve survival in acute respiratory distress syndrome (ARDS).¹

During the coronavirus-19 (COVID-19) pandemic, a huge proportion of patients developed ARDS under mechanical ventilation (MV). In this context, epidemiological studies informed higher rates of PP use; nonetheless, the number of PP sessions reported has been similar to previous studies.^{2,3}

The inability to sustain the oxygenation improvement when back to supine position, the heavy workload imposed by proning and the potential adverse effects associated with PP might lead to discussions about the usefulness of successive PP sessions in advanced phases of late ARDS.

In this retrospective, case-series study, we describe the results of patients older than 18 years old under MV that developed moderate-severe ARDS induced by COVID-19 and required 8 or more PP sessions. The number of sessions considered for inclusion was based on the number of cycles above the first standard deviation of the PROSEVA trial (4 ± 4) and the amount of sessions commonly used in previous studies.¹⁻³ The implementation and detention protocol of PP was based on PROSEVA trial recommendations.¹

Data are expressed as mean (SD), median (IQR) and number (percentage), as appropriated. The Shapiro–Wilk test was used to test normality. The analysis of data was performed using Prism version 8.4 for Mac OS (GraphPad Software, San Diego, CA, USA) and Statal Software R. All reported P values are two-sided, and have not been adjusted for multiple comparisons. A *p* value of less than 0.05 was considered to indicate statistical significance. The study was approved by the local review board (approbation code: 10-2021). All the patients signed the informed consent.

Between 1 March 2020 and 31 July 2021, one-hundred fifty one patients were admitted with diagnosis of COVID 19 and required MV. One-hundred and six (70%) required PP and twenty three (17.6%) of them needed ≥ 8 PP sessions. Demographic characteristics and outcomes of the overall sample and those who died and survived are shown in Table 1. Demographic characteristics did not significantly differ between survivors and non-survivors.

All patients met moderate to severe ARDS criteria at the time of the first PP session. A total of 273 PP sessions were performed, with a median (IQR) of 11 (9–14) sessions per patient, reaching a maximum of 21 in 1 patient.

The median (IQR) duration of invasive ventilation was 30 (23–34) days. Survivors had significantly more days of MV than non-survivors [median: 31 (IQR 30–41) vs 22 (IQR 18.5–27) $p < 0.003$]. Only 26% of the patients were weaned from MV and 60% ($n = 14$) of them underwent tracheostomy. These outcomes are representative of a subgroup of patients with infrequent characteristics in relation to the number of prone sessions reported by Estenssoro et al. and COVID-ICU Group.^{2,3} ICU mortality was 43.5%, lower than reported in mechanically ventilated patients with COVID-19 in Argentina (57%).³

Fig. 1 shows the behavior of tidal volume adjusted by predicted body weight (VT/PBW), respiratory system mechanics and oxygenation before and after the first, fifth and last PP cycle. The decision to show the fifth PP session was based on the half our PP sessions in the overall sample. All the patients presented a trend to improve the respiratory system compliance, airway driving pressure and oxygenation in the first and fifth session. This behavior was homogeneous, preventing the identification of patterns of response between survivors and non-survivors. Shelhamer et al. and Weiss et al. reports similar oxygenation outcomes in the first PP cycle.^{4,5}

Although we did not find differences in VT/PBW between groups, patients who did not survive presented higher DPaw and worst oxygenation before the last prone cycle ($p < 0.05$).

The ratio of the partial pressure of oxygen in arterial blood to the inspired oxygen fraction ($\text{PaO}_2/\text{FiO}_2$) value after the last prone session was significantly higher in survivors compared to non-survivors ($p = 0.001$). Also, the magnitude of change in $\text{PaO}_2/\text{FiO}_2$ after the last cycle was significantly different between groups ($p = 0.0001$).

In the last proning session, oxygenation only increased in patients that survived. Such finding should not be considered as a distinctive feature between groups or a predictor of results but probably reflects a clinical improvement in survivors that took us to stop proning them while an unfavorable evolution of non-survivors. We believe that the lack of improvement in gas exchange should not be considered as a stop-proning criteria.⁶ Regarding oxygenation criteria, the PROSEVA trial proposed to stop prone treatment only when $\text{PaO}_2/\text{FiO}_2$ decreased $> 20\%$ in two consecutive sessions, a situation that was not achieved by none of non-survivors. This is supported by the brief period of time from the last prone session to the end of mechanical ventilation in non-survivors (1.5 [IQR 0–2]), reinforcing that prone treatment was maintained until the last days (Table 1). In contrast to our results, Gleissman et al. found a significant increase in oxygenation only during the first three PP cycles, explaining that the lack of improvement could be due to progression of ARDS to a more proliferative phase after several days of MV. According to this observation and the high rates of complications reported during PP maneuvers, the authors suggest that assuming PP at this time point implies more risks than benefits.⁷ In our cohort, only four potentially serious complications were registered during 273 maneuvers (hypotension, oxygen desaturation and ventilator circuit disconnection) and none of the cycles had to be interrupted due to severe complications. Pressure sores (PS) were registered in 13/23 patients (56%), which were localized in the facial (10/23) and thoracic region (3/23) with a median of 2.3 (IQR 2–3) per patient. The median (IQR) grade of PS was 2 (1–2).⁸

Table 1 Patient’s characteristics and outcomes.

	All patients (n = 23)	Survivors (n = 13)	Non-survivors (n = 10)	P value
Demographic and laboratory				
Male gender, n (%)	12 (52.2)	8 (61.5)	4 (40)	0.546
Age, mean (SD)	58.2 (13.5)	55.4 (13.9)	61.8 (17.9)	0.266
SAPS II, median [IQR]	31.5 [26.5–36.5]	31 [27–38]	33 [25–35.5]	0.968
APACHE II, mean (SD)	12.6 (5.2)	12.8 (4.9)	12.1 (35.8)	0.805
Days of symptoms onset to intubation, median IQR	8 [6–11.5]	9 [8–13]	7.5 [6–8.7]	0.152
Prone positioning				
PaO ₂ /FiO ₂ before 1st PP session, mean (SD)	120.1 (25.1)	119.1 (27.3)	121.6 (23.3)	0.830
Crs before 1st PP, mean (SD)	29.4 (12)	29.3 (13.3)	30.5 (10.4)	0.831
Ventilatory Ratio before 1st PP, mean (SD)	2.2 (0.6)	2.2 (0.6)	2.2 (0.6)	0.928
PP sessions, median [IQR]	11 (9-14)	10 (9-14)	11 (9-14)	0.570
Days from last PP to end of MV, mean (SD)	7 [2–15]	13 [10–15]	1.5 [0–2]	<0.001
Outcomes				
MV days, median [IQR]	30 [23–34]	31 [30–41]	22 [18.5–27.7]	0.003
Extubation, n (%)	3 (13)	1 (7.7)	2 (20)	0.807
Reintubation, n (%)	2 (66)	1 (100)	1 (50)	1.000
Tracheostomy, n (%)	14 (60)	12 (92.3)	2 (20)	0.002

References. SAPS II: Simplified Acute Physiology Score II, APACHE II: Acute Physiology and Chronic Health Evaluation II, PaO₂/FiO₂: ratio of the partial pressure of oxygen in arterial blood (PaO₂) to the inspired oxygen fraction (FiO₂), MV: mechanical ventilation, Crs: Respiratory system compliance, PP: prone position; SD: standard deviation; IQR: interquartile range. Values are expressed in mean (SD), median [IQR] and number (percentage), as appropriate.




Figure 1 PaO₂/FiO₂ ratio, respiratory system compliance, DPaw and VT/PBW before and after the first, fifth and last prone position cycle. PaO₂/FiO₂: ratio of the partial pressure of oxygen in arterial blood (PaO₂) to the inspired oxygen fraction (FiO₂); Crs: Respiratory system compliance; DPaw: airway driving pressure; VT/PBW: tidal volume to predicted body weight; S-1: Supine before first session; P-1: after first prone session; S-5: Supine before fifth prone session; P-5: After fifth prone session; S-last: Supine before last prone session; P-last: after last prone session. Circles and squares correspond to the mean and mustaches to the standard deviations. **p* = 0.01: comparison of DPaw between survivors and non-survivors in S-last. ϕ *p* = 0.0001: comparison of the magnitude of change in PaO₂/FiO₂: between survivors and non-survivors in supine vs prone position. **p* = 0.001: comparison of PaO₂/FiO₂: between survivors and non-survivors in S-last and P-last.

Our findings support the proposed detention criteria of PROSEVA trial.¹ The favorable response even after multiple PP sessions, the low rate of severe adverse effects, the favorable results in terms of mortality and the lack of predictors associated with degree of response to the intervention suggest that interrupting the implementation of PP should not be considered even beyond the number of cycles commonly reported in the literature.

Funding sources

This research did not receive any specific grant.

Conflict of interest

None.

Acknowledgments

Lic. Daniela Gilgado (Physical and Respiratory Therapy Service Sanatorio Anchorena San Martín)

Dr. Diego Halac (Chief of Medical Clinic Sanatorio Anchorena San Martín).

References

- Guérin C, Reignier J, Richard JC, Beuret P, Gacouin A, Boulain T, et al. Prone positioning in severe acute respiratory distress syndrome. *N Engl J Med*, 368:2159–68. <https://doi.org/10.1056/NEJMoa1214103>.
- COVID-ICU Group on behalf of the REVA Network and the COVID-ICU Investigators. Clinical characteristics and day-90 outcomes of 4244 critically ill adults with COVID-19: a prospective cohort study. *Intensive care medicine*, 47:60–73. <https://doi.org/10.1007/s00134-020-06294-x>.
- Estenssoro E, Loudet CI, Ríos FG, Kanoore Edul VS, Plotnikow G, Andrian M, et al. Clinical characteristics and outcomes of invasively ventilated patients with COVID-19 in Argentina (SATICOVID): a prospective, multicentre cohort study. *Lancet Respir Med* 9:989–98. [https://doi.org/10.1016/S2213-2600\(21\)00229-0](https://doi.org/10.1016/S2213-2600(21)00229-0).
- Shelhamer MC, Wesson PD, Solari IL, Jensen DL, Steele WA, Dimitrov VG, et al. Prone positioning in moderate to severe acute respiratory distress syndrome due to COVID-19: a cohort study and analysis of physiology. *J Intensive Care Med* 36:241–52. <https://doi.org/10.1177/0885066620980399>.
- Weiss TT, Cerda F, Scott JB, Kaur R, Sungurlu S, Mirza SH, et al. Prone positioning for patients intubated for severe acute respiratory distress syndrome (ARDS) secondary to COVID-19: a retrospective observational cohort study. *Br J Anaesth* 126:48–55. <https://doi.org/10.1016/j.bja.2020.09.042>.
- Albert RK, Keniston A, Baboi L, Ayzac L, Guérin C. Proseva Investigators (2014). Prone position-induced improvement in gas exchange does not predict improved survival in the acute respiratory distress syndrome. *Am J Respir Crit Care Med* 189:494–6. <https://doi.org/10.1164/rccm.201311-2056LE>.
- Geissman H, Forsgren A, Andersson E, Lindqvist E, Lipka Falck A, Cronhjort M, et al. Prone positioning in mechanically ventilated patients with severe acute respiratory distress syndrome and coronavirus disease 2019. *Acta Anaesth Scand* 65:360–3. <https://doi.org/10.1111/aas.13741-7>.
- Benaim F, Neira J. Primer Consenso de Úlceras por Presión (PriCUPP). Bases para la implementación de un Programa de prevención, diagnóstico y tratamiento de las Úlceras por Presión. Academia Nacional de Medicina de Buenos Aires; 2017.

M. Accoce^{a,b,*}, M. Calvo Delfino^a, G. Cardoso^a, L. Castro^a, J. Pérez^a, J.H. Dorado^a

^a *Physical and Respiratory Therapy Service, Sanatorio Anchorena San Martín, Buenos Aires, Argentina*

^b *Facultad de Medicina y Ciencias de la Salud, Universidad Abierta Interamericana (UAI), Buenos Aires, Argentina*

* Corresponding author.

E-mail address: matiasaccoce@gmail.com (M. Accoce).

0210-5691/ © 2022 Elsevier España, S.L.U. y SEMICYUC. All rights reserved.

Contents of the reception guide for relatives of Spanish intensive care units: Multicenter study



Contenido de la guía de acogida para familiares de las unidades de cuidados intensivos españolas: Estudio multicéntrico

Information is one the main needs for the families of critically ill patients, and handing out to them a leaflet or a welcome guide at admission contributes to reducing the levels of anxiety involved.¹ Back in 2017, the Spanish Society of Intensive Care Nursing and Coronary Units (SEEIUC) pub-

lished a set of recommendations on how to welcome family members to intensive care units (ICU) recommending the use of written guidelines with information on the characteristics and rules of the ICU, aspects regarding the patient, and questions associated with the family.²

The objective of this study was to describe the content of the welcome guides of Spanish ICUs based on the recommendations set forth by the SEEIUC.

This was a descriptive, cross-sectional, and multicenter trial conducted from September 2019 through February 2020 that included 280 Spanish ICUs of adult patients based on a list created by the Spanish Society of Intensive and Critical Care Medicine and Coronary Units (SEMICYUC). The content of such recommendations was evaluated. The different welcome guides were obtained after contacting personnel from the different ICUs via e-mail, phone call or social media like Twitter or just by checking the hospital official website.

Data from 228 Spanish ICUs (81.4%) were eventually collected with representation from all the different Spanish

DOI of original article: <https://doi.org/10.1016/j.medin.2022.01.007>.