



ORIGINAL ARTICLE

Intrarater and interrater reliability of the Clinical Frailty Scale-Es and FRAIL-Es in critically ill patients

Susana Arias-Rivera^{a,b}, María Mar Sánchez-Sánchez^c, Raquel Jareño-Collado^c,
Marta Raurell-Torredà^{d,*}, Lorena Oteiza-López^c, Sonia López-Cuenca^c,
Israel John Thuissard-Vasallo^e, Fernando Frutos-Vivar^c

^a Programa de Doctorado en Enfermería y Salud, Facultad de Enfermería, Universidad de Barcelona, L'Hospitalet de Llobregat, Barcelona, Spain

^b Departamento de Investigación de Enfermería, Hospital Universitario de Getafe, Getafe, Madrid, Spain

^c Unidad de Cuidados Intensivos, Hospital Universitario de Getafe, Getafe, Madrid, Spain

^d Facultad de Enfermería, Universidad de Barcelona, L'Hospitalet de Llobregat, Barcelona, Spain

^e Departamento de Medicina, Facultad de Ciencias Biomédicas y de la Salud, Universidad Europea de Madrid, Villaviciosa de Odón, Madrid, Spain

Received 21 July 2024; accepted 17 September 2024

KEYWORDS

Adult;
Critical care;
Frailty;
Intensive care units;
Reliability and
validity;
Test reliability

Abstract

Objective: To evaluate the intrarater and interrater reliability of the Clinical Frailty Scale-Spain (CFS-España) and FRAIL-España and the internal consistency of the FRAIL-España when implemented in critically ill patients by intensive care nurses and physicians.

Design: Descriptive, observational and metric study.

Setting: intensive care unit (ICU) of Spain.

Patients: Patients >18 years, with >48 UCI hours.

Intervention: None.

Main variables of interest: On admission, frailty with CFS-España and FRAIL-España (by 3 nurses and 2 intensive care physicians), sex, age, comorbidities and severity.

Results: 1045 assessments were performed in 206 patients. Not frail patients on admission: 53% according to the CFS-Spain and 34% according to the FRAIL-Spain.

The intraclass correlation coefficient (ICC) shows almost perfect intrarater concordance (>0.80 for CFS-España and >0.90 for FRAIL-España). Agreement by frailty strata (non-fragile, pre-fragile and fragile patients) was substantial or almost perfect, with no major differences in ratings between nurses and physicians.

DOI of original article: <https://doi.org/10.1016/j.medin.2024.502131>

* Corresponding author.

E-mail address: mraurell@ub.edu (M. Raurell-Torredà).

<https://doi.org/10.1016/j.medicine.2025.502131>

2173-5727/© 2025 The Author(s). Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article as: S. Arias-Rivera, M.M. Sánchez-Sánchez, R. Jareño-Collado et al., Intrarater and interrater reliability of the Clinical Frailty Scale-Es and FRAIL-Es in critically ill patients, Medicina Intensiva, <https://doi.org/10.1016/j.medicine.2025.502131>

Interprofessional concordance shows an almost perfect ICC for both scales. The lowest agreement was obtained for the FRAIL-España ratings among physicians. In the frailty strata analysis, agreement was moderate. The highest agreement for the CFS-España was considering level 4 patients as frail. High reliability of the FRAIL-España and strong correlation of all dimensions with the global assessment were obtained, except for the comorbidities dimension, with a weak correlation.

Conclusion: The CFS-España and FRAIL-España scales are reliable for assessing frailty in critically ill patients by nurses and/or intensive care physicians.

© 2025 The Author(s). Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Adulto;
Cuidados críticos;
Fragilidad;
Unidades de cuidados intensivos;
Confiabilidad y validez;
Validez del test

Fiabilidad intraobservador e interobservador de las escalas de fragilidad *Clinical Frailty Scale*-España y FRAIL-España en pacientes críticos

Resumen

Objetivo: Evaluar la fiabilidad de la Clinical Frailty Scale-España (CFS-España) y la consistencia interna y fiabilidad de la FRAIL-España, implementadas en pacientes críticos por enfermeras/os de intensivos e intensivistas.

Diseño: Estudio descriptivo, observacional de carácter métrico.

Ámbito: Unidad de cuidados intensivos (UCI) de España.

Pacientes: Pacientes >18 años, con estancia en UCI > 48 horas.

Intervención: Ninguna.

Variables de interés principales: Al ingreso, fragilidad con CFS-España y FRAIL-España (por 3 enfermeras y 2 intensivistas), sexo, edad, comorbilidades (Charlson) y gravedad (SAPS3).

Resultados: Se realizaron 1045 valoraciones a 206 pacientes. Pacientes no frágiles al ingreso: 53% con CFS-España y 34% con FRAIL-España.

El Coeficiente de Correlación Intraclase (CCI) muestra una concordancia intraobservador casi perfecta (>0,80 la CFS-España y >0,90 la FRAIL-España). El acuerdo por estratos de fragilidad (pacientes no frágiles, prefrágiles y frágiles) fue sustancial o casi perfecto, sin grandes diferencias en las valoraciones entre enfermeras e intensivistas.

La concordancia entre profesionales muestra un CCI casi perfecto para ambas escalas. La concordancia más baja se obtuvo en las valoraciones de la FRAIL-España entre intensivistas. Analizando por estratos de fragilidad, el acuerdo fue moderado. El mayor acuerdo de la CFS-España fue considerando a los pacientes del nivel 4 como frágiles.

Se ha obtenido alta confiabilidad de la FRAIL-España y fuerte correlación de todas las dimensiones con la valoración global, excepto la dimensión de comorbilidades, con correlación débil. **Conclusiones:** Las escalas CFS-España y FRAIL-España son fiables para ser utilizadas en la valoración de la fragilidad de pacientes críticos aplicadas por enfermeras/os de intensivos e intensivistas.

© 2025 The Author(s). Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Frailty, understood as an increased vulnerability to unexpected outcomes from seemingly minor events (e.g., a new drug, an infection, or a minor surgical procedure), can lead to unfavorable clinical progress.¹ Frail patients are at greater risk of adverse events during hospitalization² and, upon admission to an intensive care unit (ICU), are more likely to develop post-ICU syndrome,³ experience worsening frailty,^{4,5} and face increased dependency at hospital discharge.⁶

Although there are numerous scales to evaluate frailty,⁷ none is considered as the gold standard. Two commonly used scales, likely due to their simplicity,

are the Clinical Frailty Scale (CFS)^{8,9} and the FRAIL scale.¹⁰

The Clinical Frailty Scale (CFS), designed by Rockwood et al.,¹¹ is a 9-level tool that assesses physical fitness through exercise and dependency, categorizing patients as non-frail (levels 1–3), vulnerable (level 4), and frail (mild, moderate, severe, or very severe frailty; levels 5–8). Patients at level 9 are those with a life expectancy < 6 months, regardless of evident signs of frailty. Additionally, the scale considers all patients with dementia as frail, categorizing them into mild, moderate, or severe frailty based on the degree of dementia and dependency.

The FRAIL scale^{10,12} evaluates the presence or absence of 5 dimensions—Fatigue, Resistance, Ambu-

lation, illness, and unintentional weight loss—which form the acronym. Each dimension present adds 1 point, resulting in a score between 0 and 5. Patients are categorized as non-frail (FRAIL=0), pre-frail (FRAIL=1–2), or frail (FRAIL=3–5).

In Spain, these scales are available in Spanish-adapted versions (CFS-España¹³ and FRAIL-España¹⁴) and have been applied to cohorts of critical patients.

In a study conducted in 4 Spanish ICUs,¹⁵ the prevalence of frailty—evaluated using the FRAIL scale—among patients aged 65 or older was 34.9%, which was associated with mortality 1 and 6 months after ICU discharge. In a systematic review of ICU studies,¹⁶ where frailty was measured using various tools—predominantly the CFS—the prevalence was 30%, also correlated with higher in-hospital mortality rates and long-term mortality. Identifying frail individuals is essential to help healthcare professionals individualize therapeutic limits,^{17,18} adapt care plans, and provide individualized care^{19,20} to prevent adverse events. This requires the use of valid and reliable tools to assess frailty.

Several authors have evaluated the reliability of the CFS in various contexts^{21–33} with mixed results. Published intraclass correlation coefficients (ICCs) for the CFS range from 0.350²⁹ up to 0.902⁶ for inter-observer reliability and from 0.862²² up to 0.999³⁰ for the intra-observer one. Regarding the FRAIL scale, intra-observer reliability (ICC) ranges from 0.708³⁴ up to 0.823,³⁵ with internal consistency—Kuder-Richardson Formula 20—from 0.447³⁶ up to 0.53.³⁷

The aim of this study was to evaluate the intra- and inter-observer reliability of the Clinical Frailty Scale-España, as well as the internal consistency and intra- and inter-observer reliability of the FRAIL-España when implemented in critical care patients by ICU nurses and intensivists.

Patients and methods

Study design

We conducted a descriptive, observational, metric study in a polyvalent ICU of a teaching hospital. The Guidelines for Reporting Reliability and Agreement Studies (GRRAS) checklist was followed.³⁸

Population and sample

Patients >18 years of age with expected ICU stays > 48h who consented to participate and were admitted from January 2020 through June 2023 were included. Patients with suspected imminent death or COVID-19 were excluded.

Based on the COSMIN design checklist for outcome measurement instruments,³⁹ at least, 100 pairs of evaluations by 2 independent observers would be required to analyze the inter-observer reliability. Considering a 10% dropout rate, a minimum of 111 evaluation pairs was needed.

Variables

Upon admission, frailty was assessed using the Clinical Frailty Scale-España (CFS-Es)¹³ and the FRAIL-España (FRAIL-Es).¹⁴ Other recorded variables included sex, age, comorbidities (Charlson Comorbidity Index⁴⁰), and severity level (Simplified Acute Physiology Score 3 [SAPS 3]⁴¹).

Reliability and internal consistency assessment

All participants in the study were assessed for frailty based on their condition in the month prior to hospital admission, using the CFS-Es and FRAIL-Es. The evaluations were conducted in person by 3 intensive care nurses (N1, N2, and N3), with more than 20 years of ICU experience each, and 2 intensivists (I1 and I2) with more than 10 years of specialty experience. Assessments were performed with patients directly or their relatives when patients were not capable of communicating. Not all patients could be evaluated multiple times or by all evaluators. One intensivist (I1) conducted only 1 single evaluation (always with the patient), while the remaining professionals conducted up to 2 evaluations (with relatives and/or the patient), at least, 72 h after the first evaluation. All evaluations were conducted independently. If multiple evaluators coincided during a patient visit, all could ask questions, but the individual frailty scores assigned were always blinded to the other evaluators. Upon patient discharge, the lead investigator had access to each individual evaluation from each evaluator. Although the professionals conducting the frailty evaluations were familiar with the scales, the use of such scales was not part of their routine clinical practice.

The evaluations by N1 were considered the reference among nurses because she was the study lead investigator and conducted the initial interview with the patients or their relatives, which also included questions about dependency and quality of life. Additionally, she was the only nurse to assess all patients included in the study. Evaluations by I2 were considered the reference among intensivists, as he was the only one who conducted 2 evaluations per patient. Comparisons across nurses were established between N1 and N3 due to the smaller number of assessments conducted by N2.

Data analysis

Quantitative variables are expressed as median and interquartile range [Q1–Q3], following non-parametric testing (Shapiro-Wilk). Qualitative variables are expressed as absolute (n) and relative (%) frequencies. Group median comparisons were performed using the Mann-Whitney *U* test.

Concordance was evaluated both within the same professional and among 2 and 3 professionals for the CFS-Es and FRAIL-Es scales using the ICC, through the comparison of variances (ANOVA) with repeated measures,⁴² when the scales were considered as quantitative variables. Cohen's Kappa (K)⁴³ coefficient was used for comparisons between 2 observers, and Fleiss' Kappa⁴⁴ was used to assess concordance among 3 observers. Concordance was classified as slight if ≤ 0.20 , fair if 0.21–0.40, moderate if between 0.41

Table 1 Stratifications of the Clinical Frailty Scale-Spain (CFS-Es) and the FRAIL-España (FRAIL-Es).

CFS-Es	Non-frail	Vulnerable	Frail
CFS-Es (a)	1–3	4	5–9
CFS-Es (b)	1–4	–	5–9
CFS-Es (c)	1–3	–	4–9
FRAIL-Es	Non-frail	Pre-frail	Frail
FRAIL-Es (a)	0	1–2	3–5
FRAIL-Es (b)	0–2	–	3–5

CFS-Es: Clinical Frailty Scale-Spain; FRAIL-Es: FRAIL-España.

and 0.60, substantial if between 0.61 and 0.80, and almost perfect if > 0.80.

Frailty assessments using the CFS-Es stratified patients into 3 options: option A categorized patients into 3 groups: non-frail (CFS-Es = 1–3); vulnerable or with very mild frailty (CFS-Es = 4); frail (CFS-Es = 5–9). Options B and C categorized patients into 2 groups (non-frail and frail): In option B, patients at level 4 were considered non-frail (frail = CFS-Es 5–9). In option C, patients at level 4 were considered frail (frail = CFS-Es 4–9). These stratifications were introduced following a change in nomenclature for level 4 in 2020,⁹ when it shifted from vulnerable (non-frail) to with very mild frailty (frail). Assessments using the FRAIL-Es stratified patients into 3 groups (option A): non-frail (FRAIL-Es = 0); pre-frail (FRAIL-Es = 1–2); or frail (FRAIL-Es = 3–5), or 2 groups (Option B): FRAIL-Es scores from 0 to 2 categorized patients as non-frail (Table 1).

Since the FRAIL-Es includes dichotomous items, internal consistency was estimated using the Kuder-Richardson Formula 20.⁴⁵ Reliability was categorized as very low (<0.20), low (0.21–0.40), moderate (0.41–0.60), high (0.61–0.80), and very high: (>0.80).⁴⁶ The correlation of each item with the overall scale score was evaluated using the Spearman correlation coefficient and was categorized as null (<0.10), weak (0.10–0.29), moderate (0.30–0.50), and strong correlation (>0.50).⁴⁷ The internal consistency of the FRAIL-Es was calculated using all evaluations conducted by the 5 professionals.

All values are presented with 95% confidence intervals (CI95%) and p-values, with statistical significance set at $p < 0.05$. Statistical analyses were performed using IBM SPSS Statistics for Windows (version 29.0, IBM Corp., Armonk, NY, United States).

Ethical considerations

The study protocol was reviewed and approved by the hospital Research Ethics Committee (CEIm2019/42). Consent was obtained from patients or their closest relatives when personal consent was not possible. Data confidentiality was maintained by assigning alphanumeric codes known only to the principal investigator.

Results

A total of 212 patients were included in the study, 6 of whom were excluded from this analysis because they were

Table 2 Population descriptive statistics.

	N = 206
Gender, <i>women</i> , n (%)	83 (40)
Age, <i>years</i> , <i>Median</i> [Q1–Q3]	72 [60–79]
Age, n (%)	
<50 years	27 (13)
50–65 years	48 (23)
>65 years	131 (64)
Charlson index, <i>points</i> , <i>Median</i> [Q1–Q3]	4 [3–6]
Comorbidities, n (%)	
0	47 (23)
1–2	117 (57)
>2	42 (20)
SAPS3, <i>points</i> , <i>Median</i> [Q1–Q3]	62 [53–72]
Mortality, n (%)	
In ICU	15 (7)
In hospital	32 (16)
CFS-Es, <i>level</i> , <i>Median</i> [Q1–Q3] ^a	3 [3–4]
CFS-Es, <i>level</i> , n (%) ^a	
1	9 (4)
2	24 (12)
3	76 (37)
4	62 (30)
5	19 (9)
6	11 (5)
7	5 (2)
8	0 (0)
9	0 (0)
FRAIL-Es, <i>level</i> , <i>Median</i> [Q1–Q3] ^a	1 [0–2]
FRAIL-Es, <i>dimensions present</i> , n (%) ^a	
0	70 (34)
1	48 (23)
2	39 (19)
3	32 (16)
4	14 (7)
5	3 (1)
FRAIL-Es, <i>dimensions</i> , n (%) ^a	
Fatigue (F)	91 (44)
Resistance (R)	72 (35)
Ambulation (A)	54 (26)
Illness (I)	9 (4)
Weight loss (L)	66 (32)

CFS-Es: Clinical Frailty Scale-Spain; FRAIL-Es: FRAIL-España.

^a Sourced from the first evaluation by N1 (reference professional).

only evaluated once by a single professional. A total of 1045 evaluations were conducted using each scale, consisting of 691 first evaluations (I1 = 105; I2 = 191; N1 = 206; N2 = 70; N3 = 119) and 354 s evaluations (I1 = 0; I2 = 40; N1 = 176; N2 = 43; N3 = 95).

Among the 206 included patients, 40% were women, and the median age was 72 years [60–79]. According to assessments using the CFS-Es scale (CFS-Es = 1–3), 109 (53%) of the patients were not frail at admission vs 70 (34%) according to the FRAIL-Es evaluation. The most common dimension was fatigue [91 patients (44%)], and the least common one, comorbidities [9 patients (4%)] (Table 2). The median Charlson index value was 4 [3–6], with significant differences between non-frail and frail patients

Table 3 Intra-observer reliability of the Clinical Frailty Scale-Spain and the FRAIL-Spain.

Variable	Intensivist 2 (N= 40)	Nurse 1 (N= 176)	Nurse 2 (N= 43)	Nurse 3 (N= 95)
CFS-Es, ICC (95%CI)	0.951 (0.909–0.974)	0.968 (0.957–0.976)	0.844 (0.714–0.916)	0.978 (0.967–0.985)
Kappa (95%CI); agreement (%)				
CFS-Es (a)	0.784 (0.610–0.958); 88%	0.827 (0.717–0.936); 87%	0.656 (0.424–0.889); 84%	0.874 (0.720–1.000); 93%
CFS-Es (b)	0.804 (0.542–1.000); 95%	0.898 (0.717–1.000); 97%	0.726 (0.439–1.000); 95%	0.770 (0.569–0.971); 95%
CFS-Es (c)	0.848 (0.682–1.000); 93%	0.863 (0.715–1.000); 93%	0.624 (0.329–0.919); 79%	0.936 (0.736–1.000); 97%
FRAIL-Es, ICC (95%CI)	0.902 (0.815–0.948)	0.908 (0.876–0.931)	0.924 (0.860–0.958)	0.966 (0.950–0.978)
Kappa (95%CI); agreement (%)				
FRAIL-Es (a)	0.605 (0.399–0.811); 75%	0.686 (0.580–0.792); 80%	0.858 (0.646–1.000); 91%	0.872 (0.729–1.000); 92%
FRAIL-Es (b)	0.717 (0.465–0.969); 90%	0.638 (0.490–0.785); 87%	0.743 (0.457–1.000); 91%	0.891 (0.690–1.000); 96%
Fatigue (F)	0.651 (0.353–0.948); 85%	0.730 (0.584–0.877); 87%	0.538 (0.239–0.837); 81%	0.883 (0.682–1.000); 95%
Resistance (R)	0.581 (0.285–0.877); 83%	0.818 (0.671–0.966); 92%	0.894 (0.597–1.000); 95%	0.771 (0.571–0.972); 89%
Ambulation (A)	0.817 (0.508–1.000); 93%	0.534 (0.388–0.679); 81%	0.673 (0.381–0.964); 86%	0.881 (0.681–1.000); 95%
Diseases (I)	1.000 (0.690–1.000); 100%	1.000 (0.852–1.000); 100%	1.000 (0.701–1.000); 100%	1.000 (0.799–1.000); 100%
Weight loss (L)	0.576 (0.271–0.880); 83%	0.785 (0.638–0.932); 90%	0.683 (0.386–0.980); 86%	0.922 (0.721–1.000); 97%

ICC: intraclass correlation coefficient; 95%CI: 95% confidence interval.

CFS-Es: Clinical Frailty Scale-Spain; FRAIL-Es: FRAIL-Spain.

CFS-Es (a): non-frail (1–3), vulnerable (4), frail (5–9); CFS-Es (b): non-frail (1–4), frail (5–9); CFS-Es (c): non-frail (1–3), Frail (4–9).

FRAIL-Es (a): non-frail (0), pre-frail (1–2), frail (3–5); FRAIL-Es (b): non-frail (0–2), frail (3–5).

All comparisons $p < 0.001$.

Table 4 Inter-observer reliability of the Clinical Frailty Scale-Spain and the FRAIL-Spain.

Variable	Overall (N = 122)	Nurse vs nurse (N = 211)	Intensivist vs nurse (N = 228)	Intensivist vs intensivist (N = 96)
CFS-Es, ICC (95%CI)	0.891 (0.853–0.921)	0.901 (0.870–0.924)	0.869 (0.830–0.899)	0.838 (0.758–0.892)
Kappa (95%CI); agreement (%)				
CFS-Es (a)	0.433 (0.357–0.508); 65%	0.560 (0.444–0.676); 74%	0.493 (0.386–0.600); 68%	0.396 (0.250–0.542); 60%
CFS-Es (b)	0.401 (0.299–0.504); 83%	0.585 (0.458–0.713); 90%	0.512 (0.393–0.631); 84%	0.445 (0.239–0.650); 82%
CFS-Es (c)	0.606 (0.504–0.709); 80%	0.667 (0.544–0.790); 83%	0.651 (0.533–0.769); 83%	0.471 (0.294–0.648); 74%
FRAIL-Es, ICC (95%CI)	0.895 (0.858–0.923)	0.860 (0.817–0.894)	0.873 (0.835–0.902)	0.763 (0.645–0.842)
Kappa (95%CI); agreement (%)				
FRAIL-Es (a)	0.513 (0.440–0.586); 68%	0.511 (0.400–0.622); 68%	0.514 (0.407–0.621); 68%	0.368 (0.218–0.518); 58%
FRAIL-Es (b)	0.535 (0.433–0.638); 83%	0.597 (0.473–0.721); 84%	0.584 (0.465–0.703); 84%	0.445 (0.239–0.650); 79%
Fatigue (F)	0.415 (0.312–0.519); 72%	0.389 (0.276–0.502); 71%	0.485 (0.373–0.598); 75%	0.360 (0.177–0.542); 68%
Resistance (R)	0.491 (0.388–0.595); 78%	0.579 (0.458–0.700); 81%	0.419 (0.308–0.531); 74%	0.475 (0.312–0.637); 74%
Ambulation (A)	0.571 (0.467–0.674); 82%	0.670 (0.545–0.795); 86%	0.503 (0.388–0.618); 79%	0.438 (0.253–0.624); 76%
Diseases (I)	1.000 (0.896–1.000); 100%	1.000 (0.865–1.000); 100%	0.912 (0.783–1.000); 99%	0.492 (0.108–1.000); 98%
Weight loss (L)	0.569 (0.466–0.673); 83%	0.680 (0.555–0.805); 86%	0.585 (0.467–0.703); 83%	0.245 (0.034–0.456); 73%*

ICC: intraclass correlation coefficient; 95%CI: 95% confidence interval.

CFS-Es: Clinical Frailty Scale-Spain; FRAIL-Es: FRAIL-Spain.

CFS-Es (a): non-frail (1–3), vulnerable (4), frail (5–9); CFS-Es (b): non-frail (1–4), frail (5–9); CFS-Es (c): non-frail (1–3), frail (4–9).

FRAIL-Es (a): non-frail (0), pre-frail (1–2), frail (3–5); FRAIL-Es (b): non-frail (0–2), frail (3–5).

* $p = 0.012$; other comparisons $p < 0.001$.

Table 5 Internal consistency of the FRAIL-Spain.

	Spearman (95%CI)	p-Value
Fatigue	0.721 (0.691–0.749)	<0.001
Resistance	0.751 (0.723–0.776)	<0.001
Ambulation	0.746 (0.718–0.772)	<0.001
Diseases	0.282 (0.225–0.337)	<0.001
Weight loss	0.551 (0.507–0.592)	<0.001

95%CI: 95% confidence interval.

evaluated with the FRAIL-Es scale [Charlson, FRAIL-Es 0–2 vs 3–5; 4 [2–6] vs 5 [4–8]; $p=0.001$]. Non-frail patients also exhibited lower severity at admission vs frail patients [SAPS3, FRAIL-Es 0–2 vs 3–5; 60 [50–69] vs 68 [60–77]; $p=0.001$].

Intra-observer reliability

The ICC demonstrated near-perfect agreement between first and second evaluations by each professional, with values >0.80 for the CFS-Es and >0.90 for the FRAIL-Es (Table 3).

When stratifying the scales, agreement levels ranged from substantial to near-perfect for both the CFS-Es and FRAIL-Es, depending on the stratification options. No significant differences in reliability were seen between nurses and intensivists. The FRAIL-Es dimensions with the lowest agreement were fatigue and ambulation, while comorbidities showed the highest agreement (Table 3).

Inter-observer reliability

Agreement among different professionals, measured by ICC (95%CI), was nearly perfect for both CFS-Es and FRAIL-Es. The lowest agreement was seen in FRAIL-Es evaluations among intensivists [ICC (95%CI), 0.763 (0.645–0.842); $p < 0.001$] (Table 4).

Stratified scale analysis revealed moderate to substantial agreement for both the CFS-Es and FRAIL-Es, depending on stratification. The highest agreement for CFS-Es occurred when patients were categorized into 2 strata (non-frail and frail), considering patients at level 4 as frail (Option C). Intensivists showed lower agreement vs nurses. Also, agreement was lower between intensivists and nurses. FRAIL-Es dimensions with the lowest agreement among professionals were weight loss and fatigue, while comorbidities showed the highest agreement (Table 4).

Internal consistency of the FRAIL-Es

The FRAIL-Es showed high reliability measured by Kuder-Richardson Formula 20 (95%CI), 0.643 (0.608–0.677). Strong correlations were observed between 4 different dimensions (Fatigue, Resistance, Ambulation, and Involuntary Weight Loss) and the global scale score, while the comorbidities dimension showed weak correlation (Table 5).

Discussion

The reliability of the CFS-Es and FRAIL-Es frailty scales is high, and the FRAIL-Es demonstrates strong internal consistency, regardless of the professional implementing them.

Intra-observer reliability

The CFS-Es intra-observer reliability is consistent with the results reported by Abraham et al. in the validation of the French version implemented by nurses [ICC (95%CI), 0.87 (0.76–0.93)] and intensivists [ICC (95%CI), 0.86 (0.72–0.93)]. Other validated versions in non-critical patients also show excellent intra-observer reliability, such as the Portuguese³⁰ [ICC (95%CI), 0.999 (0.998–0.999)] and Greek versions³¹ [ICC (95%CI), 0.89 (0.85–0.92)]. The Brazilian version²⁹ reported lower reliability (ICC, 0.641). These findings confirm that the CFS-Es has very good intra-observer reliability, whether implemented by nurses or intensivists.

Furthermore, the FRAIL-Es also shows very high intra-observer reliability, outperforming the Mexican (ICC, 0.82) and Chinese (ICC, 0.708) versions. The comorbidities dimension achieved 100% agreement, likely due to its objective assessment based on patient health records, whereas other dimensions relied on information from patients or families.

Inter-observer reliability

The reliability of the CFS-Es across different observers was similar to the one reported by Vrettos et al.³¹ in their validation study of the Greek version in non-critical hospitalized elderly patients [ICC (95%CI), 0.87(0.82–0.90)]. On the other hand, the reliability data of the CFS-Es exceed that reported by Abraham et al.²² in the validation of the French version among intensivists [ICC (95%CI), 0.76(0.57–0.87)] intensive care nurses [ICC (95%CI), 0.76(0.57–0.87)] or between intensivists and nurses [ICC (95%CI), 0.75(0.56–0.87) and 0.73(0.52–0.85)], respectively, as well as the reliability data reported by Rodrigues et al.²⁹ (ICC, 0.350) in the validation of the Brazilian version among volunteers. In their validation of the Danish version, Nissen et al.²⁶ achieved higher reliability among intensivists [ICC (95%CI), 0.90(0.82–0.96)], but this validation was not conducted with patient or family interviews. Instead, it involved the evaluation of written case reports, thereby excluding potential variability in patient and family responses. Even so, their reliability data are very close to those reported for the CFS-Es.

Although we did not find any studies evaluating the reliability of the FRAIL scale in critical patients, 2 studies that evaluated it among non-hospitalized adults report reliability scores lower than those of the FRAIL-Es (ICC, 0.82 in the Mexican version³⁵ and ICC, 0.708 in the Chinese version³).⁴

Regarding agreement among groups, 5 studies implemented in critical patients evaluate the reliability of the CFS.^{23–25,28,32} In the study by Flaatten et al.,³² intensivists and intensive care nurses assessed the frailty of elderly

patients (<80 years) and categorized them into 3 strata (CFS = 1–3 not frail, CFS = 4 vulnerable, and CFS = 5–9 frail). The reliability obtained among intensivists was higher than that of the CFS-Es [K(95%CI, 0.80 (0.77–0.84)] as was the reliability among nurses [K(95%CI, 0.77(0.71–0.83)], and between intensivists and nurses [K(95%CI, 0.77(0.71–0.83) and 0.80(0.77–0.84), respectively]. This superiority may be related to the fact that it was a multicenter study with 3920 patients from 22 European countries that obtained a total of 1923 pairs of evaluations. Surkan et al.²⁸ implemented the CFS (the original English version) in 158 ICU patients older than 18 years. The reliability between an intensivist and a geriatric resident or geriatrician was much lower than that obtained for the CFS-Es [K (95%CI, 0.32 (0.17–0.46) and 0.29 (–0.11 to 0.69), respectively].

In the study by Pugh et al.,²⁵ the CFS (original in English) was also implemented by intensivists and intensive care nurses. However, in this case, patients were categorized into 2 groups (CFS = 1–4 and CFS = 5–9). The reliability obtained was more similar to that of the CFS-Es [K(95%CI, intensivist-nurse 0.59 (0.44–0.75), nurse-nurse 0.63 (0.45–0.82)], except among intensivists, in whom it was higher [K (95%CI, 0.70 (0.67–0.80)]. However, the agreement percentage among intensivists was 63% vs 82% for the CFS-Es. Similarly, Hope et al.,²⁴ who analyzed variability among researchers, patients, or relatives, obtained reliability scores among the 3 groups (CFS = 1–3, CFS = 4, and CFS = 5–9; K (95%CI, 0.43 (0.42–0.46)] similar to the overall reliability of the CFS-Es and slightly higher [K (95%CI, 0.62 (0.53 to 0.70)] when analyzing variability by categorizing patients into 2 groups (CFS = 1–4 and CFS = 5–9). In a retrospective study, Darvall et al.²³ reported a Kappa coefficient of 0.67 and an agreement percentage of 45% between the observations made by a researcher and a resident intensivist, which is a Kappa coefficient slightly higher than the CFS-Es with a lower agreement percentage. Moreover, it is important to note that the evaluations from study by Darvall et al.²³ were conducted after reviewing the patients' health records, excluding potential variability from patient or family interviews.

Finally, the inter-observer reliability by groups for the CFS-Es is lower than that reported for the Turkish²⁷ (K, 0.811) and Chinese³³ (K, 0.60) versions, both implemented in ambulatory patients older than 65 years.

We did not find any studies evaluating inter-observer agreement for the FRAIL scale or its different versions. The overall data for the FRAIL-Es are moderate, whether analyzing agreement between 2 (non-frail and frail) or 3 groups (non-frail, pre-frail, and frail). The evaluations conducted by intensivists yielded the lowest scores, likely due to moderate agreement in the illness and weight loss dimensions. The Kappa coefficient of 0.492 among intensivists in the illness dimension, was undoubtedly influenced by the sample, as the agreement among intensivists was 98%, and the 2 evaluations were based on the same health record. Unintentional weight loss showed fair agreement among intensivists and moderate or substantial agreement in other comparisons (intensivist vs nurse and nurse vs nurse, respectively), possibly due to the difficulty in quantifying this dimension. Some patients were never weighed and unaware of their weight change, though repeated questioning could lead to reflection and greater clarity in the second evaluation.

Internal consistency of FRAIL-España

The internal consistency of FRAIL-Es, measured using the Kuder-Richardson Formula 20, is higher than that observed by Dong et al.³⁴ in the Chinese version of the FRAIL (KR-20 = 0.485) and by Aprahamian et al.^{36,37} in the Brazilian Portuguese version (KR-20 (95%CI), 0.447 (0.290–0.605) and KR-20 = 0.53). Although a high internal consistency was achieved, it may not have been higher due to the low number of items (5) included in the scale.⁴⁶ The correlation between the items of the scale and the overall assessment was strong, except for the comorbidities item. This issue has also been reported by Rosas-Carrasco et al.³⁵ in the Mexican Spanish version and by Susanto et al.,⁴⁸ who obtained lower correlations than those of FRAIL-Es. Rosas-Carrasco et al.³⁵ suggested that the reason for this low correlation may be that comorbidities are not as closely related to the development of frailty as the other dimensions of the scale are. However, in our cohort, we observed that frail patients (FRAIL-Es = 3–5) had a significantly higher Charlson Index vs non-frail patients (FRAIL-Es = 0–2). Certainly, comorbidities are related to frailty, though possibly to a lesser degree than the other dimensions evaluated. Alternatively, this may be related to representativeness, as only 4% of patients exhibited the comorbidities dimension.

Strengths and limitations of the study

The primary strength of this study is evaluating whether the scales are equally reliable when implemented by intensive care nurses or intensivists. Frailty assessment should be interprofessional, as a patient's frailty level may impact nursing care and/or medical treatment.

Disagreements when evaluating the same patient (either between different evaluators or by the same evaluator at different times) could have been influenced by varying responses from patients and families. It was not always possible to conduct all evaluations with the patient or their relatives. Additionally, responses could be influenced by social desirability bias, where participants only give responses they believe to be appropriate rather than truthful. When multiple relatives were present, efforts were made to include everyone in the responses to reach a consensus, but this was not always feasible. Obtaining an objective assessment is challenging, both with a scale using closed questions (like FRAIL-Es) and with open-ended scales dependent on the professional's expertise (like CFS-Es).

Furthermore, studies evaluating reliability in critical patients are scarce, with some focusing on very specific populations (>60 or >80 years) and implemented by different professionals, making comparisons with the present study difficult.

Another limitation could be the lack of prior training for professionals implementing the scales. Although they were familiar with the scales, they did not use them routinely, and the learning curve may have influenced their reliability.⁴⁹ In this regard, comparisons with other studies were not possible, as such data were not reported.

It could also be considered a limitation the fact that not all evaluators were able to assess patient frailty at 2 dif-

ferent timeframes. While desirable, the patients' clinical and/or care conditions did not always allow this.

Recommendations for future research

After establishing the reliability of the scales when implemented in critical patients, it would be interesting to analyze the reliability of these scales in a different cohort of patients or in non-hospitalized adults.

Conclusions

The CFS-Es and FRAIL-Es scales are reliable for assessing frailty in critically ill patients, whether implemented by intensive care nurses or intensivists.

CRedit authorship contribution statement

Susana Arias-Rivera: Conceptualization, Methodology, Software, Validation, Formal Analysis, Research, Resources, Data Management, Drafting, Writing – Review & Editing, Visualization, Supervision, Project Administration, Funding Acquisition. **María Mar Sánchez-Sánchez:** Validation, Research, Resources, Writing – Review & Editing. **Raquel Jareño-Collado:** Validation, Research, Resources, Writing – Review & Editing. **Marta Raurell-Torredà:** Methodology, Validation, Research, Resources, Writing – Review & Editing, Visualization. **Lorena Oteiza-López:** Validation, Research, Resources, Writing – Review & Editing. **Sonia López-Cuenca:** Validation, Research, Resources, Writing – Review & Editing. **Israel John Thuissard-Vasallo:** Validation, Formal Analysis, Research, Resources, Writing – Review & Editing. **Fernando Frutos-Vivar:** Validation, Research, Resources, Writing – Review & Editing, Visualization.

Critical review of intellectual content: all authors reviewed the final manuscript before submission for publication.

Funding

This work received support from the Spanish Ministry of Economy and Competitiveness, ISCIII-FIS, grant PI20/01231.

Declaration of competing interest

None declared.

Acknowledgments

None declared.

References

1. Clegg A, Young J, Iliffe S, Rikkert MO, Rockwood K. Frailty in elderly people. *Lancet*. 2013;381:752–62, [http://dx.doi.org/10.1016/S0140-6736\(12\)62167-9](http://dx.doi.org/10.1016/S0140-6736(12)62167-9).
2. Joseph B, Scalea T. The consequences of aging on the response to injury and critical illness. *Shock*. 2020;54:144–53, <http://dx.doi.org/10.1097/SHK.0000000000001491>.
3. Marra A, Pandharipande PP, Girard TD, Patel MB, Hughes CG, Jackson JC, et al. Co-occurrence of post-intensive care syndrome problems among 406 survivors of critical illness. *Crit Care Med*. 2018;46:1393–401, <http://dx.doi.org/10.1097/CCM.0000000000003218>.
4. Peñuelas O, Lomelí M, Del Campo-Albendea L, Toledo SI, Arellano A, Chavarría U, et al. Frailty in severe COVID-19 survivors after ICU admission. A prospective and multicenter study in Mexico. *Med Intensiva (Engl Ed)*. 2024;48:377–85, <http://dx.doi.org/10.1016/j.medine.2024.03.002>.
5. Mas Serra A. On frailty, quality of life and post-ICU syndrome. *Med Intensiva (Engl Ed)*. 2024;48:375–6, <http://dx.doi.org/10.1016/j.medine.2024.04.015>.
6. Arias-Rivera S, Sánchez-Sánchez MM, Romero de-San-Pío E, Santana-Padilla YG, Juncos-Gozaló M, Via-Clavero G, et al. Validez predictiva de la escala de fragilidad Clinical Frailty Scale-España sobre el incremento de la dependencia tras el alta hospitalaria. *Enferm Intensiva*. 2023;35:79–88, <http://dx.doi.org/10.1016/j.enfi.2023.07.003>.
7. Buta BJ, Walston JD, Godino JG, Park M, Kalyani RR, Xue Q-L, et al. Frailty assessment instruments: systematic characterization of the uses and contexts of highly-cited instruments. *Ageing Res Rev*. 2016;26:53–61, <http://dx.doi.org/10.1016/j.arr.2015.12.003>.
8. Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, McDowell I, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ*. 2005;173:489–95, <http://dx.doi.org/10.1503/cmaj.050051>.
9. Rockwood K, Theou O. Using the clinical frailty scale in allocating scarce health care resources. *Can Geriatr J*. 2020;23:210–5, <http://dx.doi.org/10.5770/cgj.23.463>.
10. Abellan van Kan G, Rolland YM, Morley JE, Vellas B. Frailty: toward a clinical definition. *J Am Med Dir Assoc*. 2008;9:71–2, <http://dx.doi.org/10.1016/j.jamda.2007.11.005>.
11. Clinical Frailty Scale. Dalhousie University. [Accessed 22 December 2023]. Available from: <https://www.dal.ca/sites/gmr/our-tools/clinical-frailty-scale.html>.
12. Abellan van Kan G, Rolland Y, Bergman H, Morley JE, Kritchevsky SB, Vellas B. The I.A.N.A Task Force on frailty assessment of older people in clinical practice. *J Nutr Health Aging*. 2008;12:29–37.
13. Arias-Rivera S, Moro-Tejedor MN, Frutos-Vivar F, Andreu-Vázquez C, Thuissard-Vasallo IJ, Sánchez-Sánchez MM, et al. Cross-Cultural Adaptation of the Clinical Frailty Scale for Critically Ill Patients in Spain and Concurrent Validity With FRAIL -Es. *Nursing Open*. 2025;12(2):e70064, <http://dx.doi.org/10.1002/nop2.70064>.
14. Arias-Rivera S, Moro-Tejedor MN, Raurell-Torredà M, Cortés-Puch I, Frutos-Vivar F, Andreu-Vázquez C, et al. Cross-cultural adaptation of the FRAIL scale for critically ill patients in Spain. *Nurs Open*. 2023;10:7703–12, <http://dx.doi.org/10.1002/nop2.2011>.
15. López Cuenca S, Oteiza López L, Lázaro Martín N, Irazabal Jaimes MM, Ibarz Villamayor M, Artigas A, et al. Frailty in patients over 65 years of age admitted to Intensive Care Units (FRAIL-ICU). *Med Intensiva*. 2019;43:395–401, <http://dx.doi.org/10.1016/j.medin.2019.01.010>.
16. Muscedere J, Waters B, Varambally A, Bagshaw SM, Boyd JG, Maslove D, et al. The impact of frailty on intensive care unit outcomes: a systematic review and meta-analysis. *Intensive Care Med*. 2017;43:1105–22, <http://dx.doi.org/10.1007/s00134-017-4867-0>.
17. Atkins CGK, Das S. Counterpoint: is it ethically permissible to use frailty scoring to determine the allocation of medical resources under conditions of scarcity in the medical ICU? *No. Chest*. 2024;166:252–4, <http://dx.doi.org/10.1016/j.chest.2024.03.014>.

18. Wilkinson DJC. Point: is it ethically permissible to use frailty scoring to determine the allocation of medical resources under conditions of scarcity in the medical ICU? Yes. *Chest*. 2024;166:250–2, <http://dx.doi.org/10.1016/j.chest.2024.03.013>.
19. Theou O, Squires E, Mallery K, Lee JS, Fay S, Goldstein J, et al. What do we know about frailty in the acute care setting? A scoping review. *BMC Geriatr*. 2018;18:139, <http://dx.doi.org/10.1186/s12877-018-0823-2>.
20. Bryden D, Jones JP, Dhesi J, Conroy S. We must consider ageing and frailty when sharing decision making in intensive care. *BMJ*. 2024;384:q116, <http://dx.doi.org/10.1136/bmj.q116>.
21. Shears M, Takaoka A, Rochweg B, Bagshaw SM, Johnstone J, Holding A, et al. Assessing frailty in the intensive care unit: a reliability and validity study. *J Crit Care*. 2018;45:197–203, <http://dx.doi.org/10.1016/j.jcrc.2018.02.004>.
22. Abraham P, Courvoisier DS, Annweiler C, Lenoir C, Milien T, Dalmaz F, et al. Validation of the clinical frailty score (CFS) in French language. *BMC Geriatr*. 2019;19:322, <http://dx.doi.org/10.1186/s12877-019-1315-8>.
23. Darvall JN, Boonstra T, Norman J, Murphy D, Bailey M, Iwashyna TJ, et al. Retrospective frailty determination in critical illness from a review of the intensive care unit clinical record. *Anaesth Intensive Care*. 2019;47:343–8, <http://dx.doi.org/10.1177/0310057X19856895>.
24. Hope AA, Munoz M, Hsieh SJ, Gong MN. Surrogates' and researchers' assessments of prehospital frailty in critically ill older adults. *Am J Crit Care*. 2019;28:117–23, <http://dx.doi.org/10.4037/ajcc2019285>.
25. Pugh RJ, Battle CE, Thorpe C, Lynch C, Williams JP, Campbell A, et al. Reliability of frailty assessment in the critically ill: a multicentre prospective observational study. *Anaesthesia*. 2019;74:758–64, <http://dx.doi.org/10.1111/anae.14596>.
26. Nissen SK, Fournaise A, Lauridsen JT, Ryg J, Nickel CH, Gudex C, et al. Cross-sectoral inter-rater reliability of the clinical frailty scale — a Danish translation and validation study. *BMC Geriatr*. 2020;20:443, <http://dx.doi.org/10.1186/s12877-020-01850-y>.
27. Özsürekli C, Balcı C, Kızırlansanoğlu MC, Çalışkan H, Tuna Doğrul R, Ayçiçek GŞ, et al. An important problem in an aging country: identifying the frailty via 9 Point Clinical Frailty Scale. *Acta Clin Belg*. 2020;75:200–4, <http://dx.doi.org/10.1080/17843286.2019.1597457>.
28. Surkan M, Rajabali N, Bagshaw SM, Wang X, Rolfson D. Interrater reliability of the clinical frailty scale by geriatrician and intensivist in patients admitted to the intensive care unit. *Can Geriatr J*. 2020;23:235–41, <http://dx.doi.org/10.5770/cgj.23.398>.
29. Rodrigues MK, Nunes Rodrigues I, Vasconcelos Gomes da Silva DJ, de S. Pinto JM, Oliveira MF. Clinical Frailty Scale: translation and cultural adaptation into the Brazilian Portuguese language. *J Frailty Aging*. 2021;10:38–43, <http://dx.doi.org/10.14283/jfa.2020.7>.
30. Pereira Pinto M, Martins S, Mesquita E, Fernandes L. European Portuguese version of the clinical frailty scale: translation, cultural adaptation and validation study. *Acta Med Port*. 2021;34:749–60, <http://dx.doi.org/10.20344/amp.14543>.
31. Vrettos I, Voukelatou P, Panayiotou S, Kyvetos A, Kalliakmanis A, Makrilakis K, et al. Validation of the revised 9-scale clinical frailty scale (CFS) in Greek language. *BMC Geriatr*. 2021;21:393, <http://dx.doi.org/10.1186/s12877-021-02318-3>.
32. Flaatten H, Guidet B, Andersen FH, Artigas A, Cecconi M, Boumendil A, et al. Reliability of the Clinical Frailty Scale in very elderly ICU patients: a prospective European study. *Ann Intensive Care*. 2021;11:22, <http://dx.doi.org/10.1186/s13613-021-00815-7>.
33. Chou Y-C, Tsou H-H, Chan D-CD, Wen C-J, Lu F-P, Lin K-P, et al. Validation of clinical frailty scale in Chinese translation. *BMC Geriatr*. 2022;22:604, <http://dx.doi.org/10.1186/s12877-022-03287-x>.
34. Dong L, Qiao X, Tian X, Liu N, Jin Y, Si H, et al. Cross-cultural adaptation and validation of the FRAIL Scale in Chinese community-dwelling older adults. *J Am Med Dir Assoc*. 2018;19:12–7, <http://dx.doi.org/10.1016/j.jamda.2017.06.011>.
35. Rosas-Carrasco O, Cruz-Arenas E, Parra-Rodríguez L, García-González AI, Contreras-González LH, Szlejf C. Cross-cultural adaptation and validation of the FRAIL Scale to assess frailty in Mexican adults. *J Am Med Dir Assoc*. 2016;17:1094–8, <http://dx.doi.org/10.1016/j.jamda.2016.07.008>.
36. Aprahamian I, Cezar NODC, Izbic R, Lin SM, Paulo DLV, Fattori A, et al. Screening for frailty with the FRAIL Scale: a comparison with the phenotype criteria. *J Am Med Dir Assoc*. 2017;18:592–6, <http://dx.doi.org/10.1016/j.jamda.2017.01.009>.
37. Aprahamian I, Lin SM, Suemoto CK, Apolinario D, Oiring De Castro Cezar N, Elmadjian SM, et al. Feasibility and factor structure of the FRAIL scale in older adults. *J Am Med Dir Assoc*. 2017;18:367.e11–8, <http://dx.doi.org/10.1016/j.jamda.2016.12.067>.
38. Kottner J, Audige L, Brorson S, Donner A, Gajewski BJ, Hróbjartsson A, et al. Guidelines for Reporting Reliability and Agreement Studies (GRRAS) were proposed. *Int J Nurs Stud*. 2011;48:661–71, <http://dx.doi.org/10.1016/j.ijnurstu.2011.01.016>.
39. Mookink LB, Prinsen CAC, Patrick DL, Alonso J, Bouter LM, de Vet HCW, et al. COSMIN Study Design checklist for patient-reported outcome measurement instruments. [Accessed 5 September 2024]. Available from: <https://www.cosmin.nl/wp-content/uploads/COSMIN-study-designing-checklist.final.pdf#>.
40. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40:373–83.
41. Metnitz PGH, Moreno RP, Almeida E, Jordan B, Bauer P, Campos RA, et al. SAPS 3—from evaluation of the patient to evaluation of the intensive care unit. Part 1: objectives, methods and cohort description. *Intensive Care Med*. 2005;31:1336–44, <http://dx.doi.org/10.1007/s00134-005-2762-6>.
42. Shrout PE, Fleiss JL. Intraclass correlations: uses in assessing rater reliability. *Psychol Bull*. 1979;86:420–8, <http://dx.doi.org/10.1037/0033-2909.86.2.420>.
43. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977;33:159–74, <http://dx.doi.org/10.2307/2529310>.
44. Enderlein G, Fleiss JL. The design and analysis of clinical experiments. Wiley, New York - Chichester - Brisbane - Toronto - Singapore 1986, 432 S. *Biom J*. 2007;30:304, <http://dx.doi.org/10.1002/bimj.4710300308>.
45. Kuder GF, Richardson MW. The theory of the estimation of test reliability. *Psychometrika*. 1937;2:151–60, <http://dx.doi.org/10.1007/BF02288391>.
46. Ruiz Bolívar C. Instrumentos y Técnicas de Investigación Educativa: Un Enfoque Cuantitativo y Cualitativo para la Recolección y Análisis de Datos. 3rd edn. Houston, Texas: DANAGA Training and Consulting; 2013.
47. Hernández Lalinde JD, Espinosa Castro F, Peñaloza Tarazona ME, Rodríguez JE, Chacón Rangel JG, Toloza Sierra CA, et al. Sobre el uso adecuado del coeficiente de correlación de Pearson: definición, propiedades y suposiciones. *Arch Venez Farmacol Terapéut*. 2018;37.

48. Susanto M, Hubbard RE, Gardiner PA. Validity and responsiveness of the FRAIL Scale in middle-aged women. *J Am Med Dir Assoc.* 2018;19:65–9, <http://dx.doi.org/10.1016/j.jamda.2017.08.003>.
49. Villa A, Poblete M. *Aprendizaje basado en competencias: una propuesta para la evaluación de las competencias genéricas [Competence-based learning: An approach to the assessment of general competences]*. Bilbao: Ediciones Mensajero; 2007.